arXiv:0903.3209v5 [math.DG] 3 Apr 2010

Uniform Approximation by Algebraic Minimal Surfaces in $\mathbb{R}^3$

Francisco J. López *

April 6, 2010

Abstract

An approximation theorem for minimal surfaces by complete minimal surfaces of finite total curvature in $\mathbb{R}^3$ is obtained. This result can be extended to the family of complete minimal surfaces of weak finite total curvature, that is to say, having finite total curvature on proper regions of finite conformal type. We deal only with the orientable case.

1 Introduction

The classical theorem of Runge deals with the uniform approximation problem for holomorphic functions on planar regions by rational functions on the complex plane. This result can be generalized to the approximation of continuous functions on Jordan curves, meromorphic functions and interpolation theorems, among other problems. Specially interesting is the approximation problem for meromorphic functions on regions of a compact Riemann surface by globally defined meromorphic functions having prescribed poles and zeroes (*algebraic* approximation). For instance, see the works by Bishop [3], Scheinberg [18, 19] and Royden [17] for a good setting.

By Huber's theorem [6], any Riemann surface M with possibly non empty compact boundary that admits a conformal complete minimal immersion with *finite total curvature* (*FTC* for short) in $\mathbb{R}^3$ has *finite conformal type*, that is to say, M is conformally equivalent to $M^c - E$, where M^c is a compact Riemann surface with possibly non empty boundary and $E \subset M^c - \partial(M^c)$ is a finite set (the topological ends of M). In addition, R. Osserman [20] proved that the Weierstrass data of the immersion extend meromorphically to the so-called Osserman compatification M^c of M. The term "algebraic" is then equivalent to the expression "of finite total curvature" in this context. Pirola [16] proved that any finitely punctured compact Riemann surface admits non-rigid conformal complete minimal immersions of FTC in $\mathbb{R}^3$. A conformal complete minimal immersion $X : M \to \mathbb{R}^3$ is said to be of *weak finite total curvature* (*WFTC* for short) if $X|_\Omega$ has FTC for all proper regions $\Omega \subset M$ of finite conformal type.

Runge's theorem has played an interesting role in the general theory of minimal surfaces. However, and contrary to its original spirit, it has taken part in very sophisticated arguments for constructing complete (or proper) minimal surfaces with arbitrary topology that are far from being algebraic in any sense (see the works of Jorge-Xavier [8], Nadirashvili [15], Lopez-Martin-Morales [13], Morales [14], Alarcon-Ferrer-Martin [2], Ferrer-Martin-Meeks [5] and Alarcón-López [1], among others).

---

*Research partially supported by MCYT-FEDER research project MTM2007-61775 and Junta de Andalucia Grant P06-FQM-01642.

2000 Mathematics Subject Classification. Primary 53A10; Secondary 49Q05, 49Q10, 53C42. Key words and phrases: Complete minimal surfaces of finite total curvature, Riemann surfaces of arbitrary conformal type, Runge's Theorem.

The aim of this paper is to obtain approximation results for complete minimal surfaces in $\mathbb{R}^3$ with FTC and *non empty compact boundary* by open complete minimal surfaces of WFTC with *arbitrarily prescribed conformal structure and flux map.* For a thorough exposition of our results, the following notations are required.

If $X : M \to \mathbb{R}^3$ is a conformal minimal immersion and $\gamma \subset M$ is an oriented closed curve, the flux of X on γ is given by $p_X(\gamma) := \int_\gamma \mu(s)ds$, where s is the oriented arclength parameter on γ and $\mu(s)$ is the conormal vector of X at $\gamma(s)$ for all s. Recall that $\mu(s)$ is the unique unit tangent vector of X at $\gamma(s)$ such that $\{dX(\gamma'(s)), \mu(s)\}$ is a positive basis. Since X is a harmonic map, $p_X(\gamma)$ depends only on the homology class of γ and the well defined flux map $p_X : H_1(M, \mathbb{Z}) \to \mathbb{R}^3$ is a group homomorphism.

As usual, a surface is said to be *open* if it is non-compact and has empty boundary.

Definition 1.1 *Let N be an open Riemann surface, and let M be a proper region of N with compact boundary. Throughout this paper, we will deal with the following spaces of minimal surfaces:*

(a) $\mathcal{E}(N)$ is the space of conformal complete minimal immersions $X : N \to \mathbb{R}^3$ of WFTC.

(b) $\mathcal{E}_N(M)$ is the space of conformal complete minimal immersions $X : M \to \mathbb{R}^3$ of WFTC that extend as a conformal minimal immersion to some neighborhood of M in N.

If M has finite conformal type, $\mathcal{E}_N(M)$ is the space of conformal complete minimal immersions of M in $\mathbb{R}^3$ with FTC that extend to some neighborhood of M in N. Obviously $\mathcal{E}(M) = \mathcal{E}_M(M)$ when M is open. All these spaces will be endowed with the $\mathcal{C}^0$ topology of *uniform convergence on (not necessarily compact) proper regions of finite conformal type in M.* To be more precise:

Definition 1.2 *A sequence $\{X_n\}_{n\in\mathbb{N}} \subset \mathcal{E}_N(M)$ is said to converge in the $\mathcal{C}^0$ topology to $X_0 \in \mathcal{E}_N(M)$ if for any proper region $\Omega \subset M$ of finite conformal type, $\{X_n|_\Omega\}_{n\in\mathbb{N}} \to X_0|_\Omega$ uniformly on Ω, that is to say, in the $\mathcal{C}^0(\Omega)$-topology associated to the norm of the supremum*[1] *on Ω. If M has finite conformal type, this topology coincides with the one of uniform convergence on M.*

Our first result is the following density theorem, in which all the involved immersions are of FTC (see Corollary 4.1):

> **Theorem I (Basic Approximation Theorem):** *Let N be an open Riemann surface of finite conformal type, let M be a compact region in N such that $N - M$ consists of a finite collection of pairwise disjoint once punctured discs and consider $X \in \mathcal{E}_N(M)$.*
>
> *Then there exists $\{Y_n\}_{n\in\mathbb{N}} \subset \mathcal{E}(N)$ such that $\{Y_n|_M\}_{n\in\mathbb{N}} \to X$ uniformly on M and $p_{Y_n} = p_X$ for all n.*

Given two Riemann surfaces M and M^* with possibly non empty compact boundary, M^* is said to be an *extension* of M if M is a proper subset of M^*, $M \cap \partial(M^*) = \emptyset$ and $M^* - M^\circ$ contains no compact connected components that are disjoint from $\partial(M^*)$, where M° is the topological interior of M in M^*. In particular the induced group homomorphism $j_* : H_1(M, \mathbb{R}) \to H_1(M^*, \mathbb{R})$ is injective, where $j : M \to M^*$ is the inclusion map (up to the natural identification we consider $H_1(M, \mathbb{R}) \subset H_1(M^*, \mathbb{R})$). This notion makes sense even when M is not connected.

We have also proved the following general approximation result (see Theorem 4.2):

> **Theorem II (General Approximation Theorem):** *Let M be a Riemann surface of finite conformal type with $\partial(M) \neq \emptyset$, and let N be any extension of M with $\partial(N) = \emptyset$. Consider $X \in \mathcal{E}_N(M)$ and a group homomorphism $q : H_1(N, \mathbb{Z}) \to \mathbb{R}^3$ such that $q|_{H_1(M,\mathbb{Z})} = p_X$.*

---

[1] By the maximum principle at infinity, in the norm of the maximum.

Then there exist $\{Y_n\}_{n\in\mathbb{N}} \subset \mathcal{E}(N)$ *such that* $\{Y_n|_M\}_{n\in\mathbb{N}} \to X$ *uniformly on* M *and* $p_{Y_n} = q$ *for all* n*.*

As a consequence, $\mathcal{E}(M^*) \neq \emptyset$ for any open Riemann surface M^*, and for any group homomorphism $q : H_1(M^*, \mathbb{Z}) \to \mathbb{R}^3$ we can find $Y \in \mathcal{E}(M^*)$ with $p_Y = q$. The key idea of the General Approximation Theorem is to prove an *algebraic bridge principle* for complete minimal immersions of FTC which allows good control over the conformal structure, asymptotic behavior and flux map (see Theorem 4.1). This bridge principle can be used in general connected sum constructions for complete minimal surfaces of finite total curvature(see [11]). Other results of this kind can be found in Kapouleas [9], Yang [23] and White [21, 22] papers.

The paper is laid out as follows. Section 2 is devoted to some preliminary results on Algebraic Geometry, and in Section 3 we go over the Weierstrass and spinorial representations of minimal surfaces. Section 4 contains the main results: the Algebraic Bridge Principle is proved in Subsection 4.1 whereas the Basic and General Approximation Theorems are proved in Subsection 4.2.

In a forthcoming paper [12] the author will extend this analysis to the nonorientable case.

2 Preliminaries on Riemann surfaces

As usual, we call $\overline{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ the extended complex plane or Riemann sphere.

Given a Riemann surface M, $\partial(M)$ will denote the one dimensional topological manifold determined by the boundary points of M, and $\mathrm{Int}(M)$ the open surface determined by the interior points of M. Obviously, $\mathrm{Int}(M) = M - \partial(M)$. Given $S \subset M$, write S° and $\overline{S}$ for the topological interior and closure of S in M, and likewise $Fr(S) = \overline{S} - S^\circ$ for the topological frontier of S. If S is a topological surface with the induced topology then $\partial(S) \subset Fr(S)$, but in general the contrary is false. A proper connected subset $S \subset M$ is said to be a *region* if it is a topological surface with the induced topology. An open connected subset of $\mathrm{Int}(M)$ will be called a *domain* of M.

Given two Riemann surfaces M and M^* with possibly non empty compact boundary, M^* is said to be an *annular extension* of M if M^* is an extension of M and $M^* - M^\circ$ consists of a finite collection of compact annuli and once punctured closed discs. If in addition $M^* - M^\circ$ is compact then M^* is said to be a *trivial* annular extension of M. In this case M and M^* are homeomorphic.

2.1 Approximation results on Riemann surfaces

Let N be a Riemann surface with $\partial(N) = \emptyset$, and let S be a subset of N with $S \neq N$. A connected component V of $N - S$ is said to be *bounded* if $\overline{V}$ is compact.

Definition 2.1 *We denote by* $\mathfrak{F}^N(S)$ *the space of continuous functions* $f : S \to \overline{\mathbb{C}}$ *such that* f *is meromorphic on some open neighborhood of* $\overline{S^\circ} \cap S$ *in* N *and* $f(P) \neq \infty$ *for all* $P \in S - \overline{S^\circ}$*. If* S *is open then* $\mathfrak{F}^N(S)$ *coincides with the space of meromorphic functions on* S*, hence does not depend on* N *and is simply written* $\mathfrak{F}(S)$*. We write* $\mathfrak{F}_0^N(S)$ *(and* $\mathfrak{F}_0(S)$ *when* S *is open) for the corresponding space of holomorphic functions.*

If V is an open subset of N and $\overline{S} \subset V$ then $\mathfrak{F}^V(S) = \mathfrak{F}^N(S)$ and $\mathfrak{F}_0^V(S) = \mathfrak{F}_0^N(S)$.

All these spaces will be endowed with the $\omega(S)$-topology, namely, the topology of the uniform convergence on S. To be more precise, we shall say that a function $f \in \mathfrak{F}^N(S)$ can be uniformly approximated on S by functions in $\mathfrak{F}(N)$ if there exists $\{f_n\}_{n\in\mathbb{N}} \subset \mathfrak{F}(N)$ such that $\{|f_n|_S - f|\}_{n\in\mathbb{N}} \to 0$ uniformly on S. In particular all f_n have the same set $\mathcal{P}_f$ of poles on S. Likewise we define the uniform approximation of an $f \in \mathfrak{F}_0^N(S)$ by functions in $\mathfrak{F}_0(N)$.

A complex 1-form θ on S is said to be of type $(1,0)$ if for any conformal chart (U, z) on N we have that $\theta|_{U\cap S} = f(z)dz$ for some $f : U \cap S \to \overline{\mathbb{C}}$. For instance, holomorphic and meromorphic 1-forms on N are of type $(1,0)$.

Definition 2.2 *We denote by $\mathfrak{W}^N(S)$ the space of 1-foms of type $(1,0)$ on S with $\theta(P) \neq \infty$ for all $P \in S - \overline{S^\circ}$ and being meromorphic on some open neighborhood of $\overline{S^\circ} \cap S$ in N. If S is open then $\mathfrak{W}^N(S)$ coincides with the space of meromorphic 1-forms on S, hence does not depend on N and is simply written $\mathfrak{W}(S)$. We write $\mathfrak{W}_0^N(S)$ (and $\mathfrak{W}_0(S)$ when S is open) for the corresponding space of holomorphic 1-forms.*

If V is an open subset of N and $\overline{S} \subset V$ then $\mathfrak{W}^V(S) = \mathfrak{W}^N(S)$ and $\mathfrak{W}_0^V(S) = \mathfrak{W}_0^N(S)$.

A 1-form $\theta \in \mathfrak{W}^N(S)$ can be uniformly approximated on S by 1-forms in $\mathfrak{W}(N)$ if there exists $\{\theta_n\}_{n\in\mathbb{N}} \subset \mathfrak{W}(N)$ such that $\{\frac{\theta_n - \theta}{dz}|_{S\cap K}\}_{n\in\mathbb{N}} \to 0$ in the $\omega(S \cap K)$-topology for any closed conformal disc (K, z) on N (we are assuming that $z : K \to z(K) \subset \mathbb{C}$ extends as a conformal parameter beyond K in N). In particular all θ_n have the same set of poles $\mathcal{P}_\theta$ on S. As above, we say that $\{\theta_n|_S\}_{n\in\mathbb{N}} \to \theta$ in the $\omega(S)$-topology. Likewise we define the uniform approximation of a $\theta \in \mathfrak{W}_0^N(S)$ by 1-forms in $\mathfrak{W}_0(N)$.

Let $\mathfrak{Div}(S)$ denote the free commutative group of divisors of S with multiplicative notation. If $D = \prod_{i=1}^n Q_i^{n_i} \in \mathfrak{Div}(S)$, where $n_i \in \mathbb{Z} - \{0\}$ for all i, the set $\{Q_1, \ldots, Q_n\}$ is said to be the *support* of D. Let $\mathfrak{Deg} : \mathfrak{Div}(S) \to \mathbb{Z}$ be the group homomorphism given by the degree map $\mathfrak{Deg}(\prod_{j=1}^t Q_j^{n_j}) = \sum_{j=1}^t n_j$. A divisor $D \in \mathfrak{Div}(S)$ is said to be *integral* if $D = \prod_{i=1}^n Q_i^{n_i}$ and $n_i \geq 0$ for all i. Given D_1, $D_2 \in \mathfrak{Div}(S)$, $D_1 \geq D_2$ if and only if $D_1 D_2^{-1}$ is integral. For any $f \in \mathfrak{F}^N(S)$ we denote by $(f)_0$ and $(f)_\infty$ its associated integral divisors of zeroes and poles in S, respectively, and label $(f) = (f)_0/(f)_\infty$ as the divisor associated to f on S. Likewise we define $(\theta)_0$ and $(\theta)_\infty$, and write $(\theta) = (\theta)_0/(\theta)_\infty$ for the divisor of θ in S, $\theta \in \mathfrak{W}^N(S)$.

E. Bishop [3], H. L. Royden [17] and S. Scheinberg [18, 19], among others, have proved several extensions of Runge's theorem. For our purposes, we need only the following compilation result:

Theorem 2.1 *Let N be a Riemann surface with $\partial(N) = \emptyset$, and let $S \subset N$ be a compact not necessarily connected subset consisting of a finite collection of pairwise disjoint compact regions in N, and a finite collection of pairwise disjoint Jordan arcs with endpoints at the boundary of these regions and otherwise disjoint from them. Let $E \subset N - S$ be a subset meeting each bounded component of $N - S$ in a unique point.*

Then any function $f \in \mathfrak{F}^N(S)$ can be uniformly approximated on S by functions $\{f_n\}_{n\in\mathbb{N}}$ in $\mathfrak{F}(N) \cap \mathfrak{F}_0(N - (E \cup \mathcal{P}_f))$, where $\mathcal{P}_f = f^{-1}(\infty) \subset S$. Furthermore, if $D \in \mathfrak{Div}(S)$ is an integral divisor, the sequence can be chosen in such a way that $(f - f_n|_S)_0 \geq D$.

2.2 Compact Riemann surfaces

The background of the following results can be found, for instance, in [4].

In the sequel, R will denote a compact Riemann surface with genus $\nu \geq 1$ and $\partial(R) = \emptyset$.

Label $H_1(R, \mathbb{Z})$ as the 1^{st} homology group with integer coefficients of R. Let $B = \{a_j, b_j\}_{j=1,\ldots,\nu}$ be a canonical homology basis of $H_1(R, \mathbb{Z})$, and write $\{\xi_j\}_{j=1,\ldots,\nu}$ the associated dual basis of $\mathfrak{W}_0(R)$, that is to say, the one satisfying that $\int_{a_k} \xi_j = \delta_{jk}, \quad j,\ k = 1, \ldots, \nu$.

Denote by $\Pi = (\pi_{jk})_{j,\,k=1,\ldots,\nu}$ the Jacobi period matrix with entries $\pi_{jk} = \int_{b_k} \xi_j, \quad j,\ k = 1, \ldots, \nu$. This matrix is symmetric and has positive definite imaginary part. We denote by $L(R)$ the lattice over $\mathbb{Z}$ generated by the 2ν-columns of the $\nu \times 2\nu$ matrix (I_ν, Π), where I_ν is the identity matrix of dimension ν.

Finally, call $J(R) = \mathbb{C}^\nu / L(R)$ the Jacobian variety of R, which is a compact, commutative, complex, ν-dimensional Lie group. Fix $P_0 \in R$, denote by $\varphi_{P_0} : \mathfrak{Div}(R) \to J(R), \quad \varphi_{P_0}(\prod_{j=1}^s Q_j^{n_j}) = \sum_{j=1}^s n_j\, {}^t(\int_{P_0}^{Q_j} \xi_1, \ldots, \int_{P_0}^{Q_j} \xi_\nu)$ the Abel-Jacobi map with base point P_0, where ${}^t(\,\cdot\,)$ means matrix transpose. If there is no room for ambiguity we simply write φ.

Abel's theorem asserts that $D \in \mathfrak{Div}(R)$ is the principal divisor associated to a meromorphic function $f \in \mathfrak{W}(R)$ if and only if $\mathfrak{Deg}(D) = 0$ and $\varphi(D) = 0$. Jacobi's theorem says that $\varphi : R_\nu \to J(R)$ is surjective and has maximal rank (hence a local biholomorphism) almost everywhere, where R_ν denotes the space of integral divisors in $\mathfrak{Div}(R)$ of degree ν.

Riemann-Roch theorem says that $r(D^{-1}) = \mathfrak{Deg}(D) - g + 1 + i(D)$ for any $D \in \mathfrak{Div}(R)$, where $r(D^{-1})$ (respectively, $i(D)$) is the dimension of the complex vectorial space of functions $f \in \mathfrak{F}(R)$ (respectively, 1-forms $\theta \in \mathfrak{W}(R)$) satisfying that $(f) \geq D^{-1}$ (respectively, $(\theta) \geq D$).

By Abel's theorem, the point $\kappa_R := \varphi((\theta)) \in J(R)$ does not depend on $\theta \in \mathfrak{W}(R)$. It is called as the *vector of the Riemann constants*. Write $\mathcal{S}(R)$ for the set containing the $2^{2\nu}$ solutions of the algebraic equation $2s = \kappa_R$ in $J(R)$. Any element of $\mathcal{S}(R)$ is said to be a *spinor structure* on R. A 1-form $\theta \in \mathfrak{W}(R)$ is said to be *spinorial* if $(\theta) = D^2$ for a divisor $D \in \mathfrak{Div}(R)$. Denote by $\mathfrak{S}(R)$ (respectively, $\mathfrak{S}_0(R)$) the set of spinorial meromorphic (respectively, spinorial holomorphic) 1-forms on R. Two 1-forms θ_1, $\theta_2 \in \mathfrak{S}(R)$ are said to be spinorially equivalent, written $\theta_1 \sim \theta_2$, if there exists $f \in \mathfrak{F}(R)$ such that $\theta_2 = f^2\theta_1$. A class $\Theta \in \frac{\mathfrak{S}(R)}{\sim}$ determines a unique spinor structure $s_\Theta \in \mathcal{S}(R)$. Indeed, it suffices to take $\theta \in \Theta$ and define $s_\Theta = \varphi(D)$, where $D^2 = (\theta)$. By Abel's theorem s_Θ does not depend on the chosen $\theta \in \Theta$.

The map $\frac{\mathfrak{S}(R)}{\sim} \to \mathcal{S}(R)$, $\Theta \mapsto s_\Theta$ is bijective. To see this, take $s \in \mathcal{S}(R)$ and use Jacobi's theorem to find an integral divisor $D' \in \mathfrak{Div}(R)$ of degree ν satisfying $\varphi(D') = s$. By Abel's theorem $(D'P_0^{-1})^2$ is the canonical divisor associated to a spinorial meromorphic 1-form whose corresponding class Θ_s in $\frac{\mathfrak{S}(R)}{\sim}$ satisfies $s_{\Theta_s} = s$.

Spinor structures can be also introduced in a more topological way. Indeed, take $s \in \mathcal{S}(R)$ and $\theta \in \Theta_s$. For any embedded loop $\gamma \subset R$, consider an open annular neighborhood A of γ and a conformal parameter $z : A \to \{z \in \mathbb{C} \ : \ 1 < |z| < r\}$. Set $\xi_s(\gamma) = 0$ if $\sqrt{\theta(z)/dz}$ has a well defined branch on A and $\xi_s(\gamma) = 1$ otherwise, and note that this number does not depend on the annular conformal chart. The induced map $\xi_s : H_1(R, \mathbb{Z}) \to \mathbb{Z}_2$ does not depend on $\theta \in \Theta_s$ and defines a group homomorphism. Furthermore, $\xi_{s_1} = \xi_{s_2}$ if and only if $s_1 = s_2$, and therefore $\mathcal{S}(R)$ can be identified with the set of group morphisms $\mathrm{Hom}(H_1(R, \mathbb{Z}), \mathbb{Z}_2)$. We also write $\xi_\Theta = \xi_{s_\Theta}$, for any $\Theta \in \frac{\mathfrak{S}(R)}{\sim}$.

2.3 Riemann surfaces of finite conformal type

Let$M$ be a Riemann surface possibly with non empty compact boundary and of finite conformal type. The *Osserman compactification* M^c of M is obtained by filling out the conformal punctures of M (that is to say, the anuular ends of M of finite conformal type). Attaching conformal discs to the boundary components of M^c, we get a compac Riemann surface R without boundary that will be called a *conformal compactification* of M. Notice that M^c is unique, whereas R depends on the gluing process of the conformal discs. With this language, $M^c = R - (\cup_{j=1}^b U_j)$ and $M = M^c - \{E_1, \ldots, E_a\}$, where U_j, $j = 1, \ldots, b$ are open discs with pairwise disjoint closures in R and $\{E_1, \ldots, E_a\} \subset M^c - \partial(M^c)$.

Let M be a Riemann surface of finite conformal type, and call $H_1(M, \mathbb{Z})$ as the 1^{st} homology group of M with integer coefficients. Set

$$\mathfrak{S}(M) = \{\theta \in \mathfrak{W}(\mathrm{Int}(M)),\ (\theta) = D^2,\ D \in \mathfrak{Div}(\mathrm{Int}(M))\}, \quad \mathfrak{S}_0(M) = \mathfrak{S}(M) \cap \mathfrak{W}_0(\mathrm{Int}(M)).$$

Two 1-forms θ_1, $\theta_2 \in \mathfrak{S}(M)$ are said to be *spinorially equivalent* if there exists $f \in \mathfrak{F}(\mathrm{Int}(M))$ such that $\theta_2 = f^2\theta_1$. Define the map $\xi : \frac{\mathfrak{S}(M)}{\sim} \to \mathrm{Hom}(H_1(M, \mathbb{Z}), \mathbb{Z}_2)$, $\Theta \mapsto \xi_\Theta$, as above.

Lemma 2.1 *The map $\xi : \frac{\mathfrak{S}(M)}{\sim} \to Hom(H_1(M, \mathbb{Z}), \mathbb{Z}_2)$ is bijective.*

Proof: Standard monodromy arguments give that ξ is injective.

Write $M = M^c - \{E_1, \ldots, E_a\}$ and $\partial(M) = \cup_{j=1}^b c_j$, where c_j is a Jordan curve for all j and $c_{j_1} \cap c_{j_2} = \emptyset$ when $j_1 \neq j_2$. Consider a family $V_1, \ldots, V_a$ of pairwise disjoint closed discs in $M^c - \partial(M)$ such that $E_i \in V_i^\circ$ for all $i = 1, \ldots, a$. Labeling $r := a + b > 0$ and $\{d_1, \ldots, d_r\} = \{\partial(V_i),\ i = 1, \ldots, a\} \cup \{c_j,\ j = 1, \ldots, b\}$, we know that $\{a_1, \ldots, a_\nu, b_1, \ldots, b_\nu, d_1, \ldots, d_{r-1}\}$ is a basis of $H_1(M, \mathbb{Z})$, so $\mathrm{Hom}(H_1(M, \mathbb{Z}), \mathbb{Z}_2)$ contains $2^{2\nu+r-1}$ elements. Let R be a conformal compatification of M, and write $\frac{\mathfrak{S}(R)}{\sim} = \{\Theta_j,\ j = 1, \ldots 2^{2\nu}\}$. Choose $\theta_j \in \Theta_j$ for each j and call $f_j = \theta_j/\theta_1 \in \mathfrak{F}(R)$, $j = 1, \ldots, 2^{2\nu}$. Since Θ_i and Θ_j correspond to different spinor structures on R, $i \neq j$, $\sqrt{\theta_i/\theta_j}$ has no well defined branches on R, hence on $\mathrm{Int}(M)$. Thus $\{\theta_j|_{\mathrm{Int}(M)}\ :\ j = 1, \ldots 2^{2\nu}\}$ are pairwise spinorially inequivalent in $\mathfrak{S}(M)$. Write $M^c = R - \cup_{j=1}^b U_j$, where U_j is an open disc in R with $\partial(U_j) = c_j$ for all j, and fix $E_{a+j} \in U_j$, $j = 1, \ldots, b$. For any $J \subseteq \{1, \ldots, r-1\}$, $J \neq \emptyset$, use Jacobi's theorem to find an integral divisor $D_J \in \mathfrak{Div}(R)$ of degree $\nu + \sharp(J) - 1$ verifying $\varphi(D_J^2 E_r^{-\sharp(J)} \prod_{j \in J} E_j^{-1}) = \kappa_R$, where $\sharp(J)$ is the cardinal of J. By Abel's theorem, there exists $\tau_J \in \mathfrak{W}(R)$ with canonical divisor $(\tau_J) = D_J^2 E_r^{-\sharp(J)} \prod_{j \in J} E_j^{-1}$. Since $f_i \tau_J/\theta_j$ has a pole of odd order at some E_h, $(f_i \tau_J)|_{\mathrm{Int}(M)}$ and $\theta_j|_{\mathrm{Int}(M)}$ are spinorially inequivalent in $\mathfrak{S}(M)$, $i, j \in \{1, \ldots, 2^{2\nu}\}$, and likewise for any pair $(f_{i_1}\tau_{J_1})|_{\mathrm{Int}(M)}$, $(f_{i_2}\tau_{J_2})|_{\mathrm{Int}(M)}$ with $(i_1, J_1) \neq (i_2, J_2)$. Thus the set $\{\theta_j|_{\mathrm{Int}(M)},\ j = 1, \ldots, 2^{2\nu}\} \cup \{(f_i \tau_J)|_{\mathrm{Int}(M)},\ i = 1, \ldots, 2^{2\nu},\ J \subseteq \{1, \ldots, r-1\},\ J \neq \emptyset\}$ consists of $2^{2\nu+r-1}$ pairwise spinorially inequivalent 1-forms in $\mathfrak{S}(M)$, proving the surjectivity of ξ. $\Box$

3 Analytic Representations of Minimal Surfaces

Let N be an open Riemann surface and $M \subset N$ a proper region with compact boundary.

Let X be a minimal immersion in $\mathcal{E}_N(M)$. Write $\partial_z X = (\phi_1, \phi_2, \phi_3)$ and notice that $\partial_z X \in \mathfrak{W}_0^N(M)^3$. Since X is conformal and minimal, then $\phi_1 = \frac{1}{2}(1/g - g)\phi_3$ and $\phi_2 = \frac{i}{2}(1/g + g)\phi_3$, where $g \in \mathfrak{F}^N(M)$ and up to the stereographic projection coincides with the Gauss map of X. The pair (g, ϕ_3) is known as the *Weierstrass representation* of X.

Clearly $X(P) = X(Q) + \mathrm{Re} \int_Q^P (\phi_1, \phi_2, \phi_3)$, P, $Q \in M$. The induced intrinsic metric ds^2 on M and its Gauss curvature $\mathcal{K}$ are given by the expressions:

$$ds^2 = \sum_{j=1}^{3} |\phi_j|^3 = \frac{1}{4}|\phi_3|^2 (\frac{1}{|g|} + |g|)^2, \quad \mathcal{K} = -\left(\frac{4|dg||g|}{|\phi_3|(1 + |g|^2)^2} \right)^2. \tag{1}$$

The total curvature of X is given by $c(X) := \int_M \mathcal{K} dA$, where dA is the area element of ds^2, and the *flux map* of X by the expression $p_X : H_1(M, \mathbb{Z}) \to \mathbb{R}^3$, $p_X(\gamma) = \mathrm{Im} \int_\gamma \partial_z X$.

By Huber, Osserman and Jorge-Meeks results [6, 20, 7], if X is complete and of FTC then X is proper, M has finite conformal type, the Weierstrass data (g, ϕ_3) of X extend meromorphically to M^c and the vectorial 1-form $\partial_z X$ has poles of order ≥ 2 at the ends.

Remark 3.1 *Recall that $\mathcal{E}_N(M)$ is endowed with the $\mathcal{C}^0$ topology of the uniform convergence on proper regions of finite conformal type (see Definition 1.2). Take $\{X_n,\ n \in \mathbb{N}\} \cup \{X\} \subset \mathcal{E}_N(M)$ and assume that $\{X_n\}_{n \in \mathbb{N}} \to X$ in $\mathcal{E}_N(M)$. If $\Omega \subset M$ is a proper region of finite conformal type, it is not hard to see that $\{(X_n - X)|_{\Omega^c}\}_{n \in \mathbb{N}} \to 0$ uniformly on Ω^c and the Weierstrass data of X_n converge in the $\omega(\Omega^c)$-topology to the ones of X on Ω^c. Indeed, just observe that by Riemann's removable singularity theorem $X_n - X$ extends harmonically to the punctures of Ω for all n, and $\{X_n - X\}_{n \in \mathbb{N}} \to 0$ uniformly on Ω^c as well.*

Assume now that M has finite conformal type, consider $X \in \mathcal{E}_N(M)$ and write (g, ϕ_3) for its Weierstrass data. Since ds^2 has no singularities on M (see equation (1)), $\eta_1 = \frac{\phi_3}{g}$ and $\eta_2 = \phi_3 g$

are 1-forms in $\mathfrak{W}_0^N(M)$ without common zeroes on M and spinorially equivalent in $\mathfrak{S}_0(M^*)$, where M^* is any trivial annular extension of M in N to which X extends. Furthermore, we know that at least one of them has a pole (of order ≥ 2) at E for each $E \in M^c - M$. The next lemma shows that the converse is true:

Lemma 3.1 (Spinorial Representation) *Let N be an open Riemann surface, let M be a proper region in N of finite conformal type and let M^* be a trivial annular extension of M in N. Let η_1, η_2 be two spinorially equivalent 1-forms in $\mathfrak{S}_0(M^*) \cap \mathfrak{W}^N(M^c)$ such that $|\eta_1| + |\eta_2|$ never vanishes in M, at least one of the 1-forms η_j, $j = 1, 2$, has a pole at E for each $E \in M^c - M$, and $\frac{1}{2}(\eta_1 - \eta_2)$, $\frac{i}{2}(\eta_1 + \eta_2)$ and $\sqrt{\eta_1 \eta_2}$ have no real periods on M.*

Then the map $X : M \to \mathbb{R}^3$,

$$X(P) = Re \int_{P_0}^{P} \left(\frac{1}{2}(\eta_1 - \eta_2), \frac{i}{2}(\eta_1 + \eta_2), \sqrt{\eta_1 \eta_2} \right), \quad P_0 \in M, \tag{2}$$

is well defined and lies in $\mathcal{E}_N(M)$.

Proof: Since η_1 and η_2 are spinorially equivalent in $\mathfrak{S}_0(M^*)$ and lie in $\mathfrak{W}^N(M^c)$, there is $g \in \mathfrak{F}^N(M^c)$ such that $\eta_2 = g^2 \eta_1$, and therefore $\phi_3 := \sqrt{\eta_1 \eta_2}$ is well defined. As $\frac{1}{2}(\eta_1 - \eta_2)$, $\frac{i}{2}(\eta_1 + \eta_2)$ and ϕ_3 have no real periods on M, then X is well defined. Furthermore, from our hypothesis $\frac{1}{4}|\phi_3|^2(\frac{1}{|g|} + |g|)^2$ never vanishes on M, hence X is the minimal immersion with Weierstrass data (g, ϕ_3). Following Osserman [20], X is complete and of FTC. $\Box$

The pair (η_1, η_2) will be called as the *spinorial representation* of X (see [10] for a good setting).

4 Approximation by complete minimal surfaces with FTC

Roughly speaking, the aim of this section is to prove that any figure in $\mathbb{R}^3$ consisting of two complete minimal surfaces with FTC and non empty compact boundary (for instance, two compact minimal surfaces) and a Jordan arc connecting them, can be uniformly approximated by complete minimal surfaces of FTC without boundary. Furthermore, the conformal structure and the flux map of the approximate sequence can be prescribed.

For a thorough explanation of these results, the following notations are required.

Let N be an open Riemann surface, and let M_1, M_2 be two proper regions in N with finite conformal type and non-empty boundary, where either $M_1 \cap M_2 = \emptyset$ or $M_1 = M_2$. Call $M = M_1 \cup M_2$ and consider an analytic regular Jordan arc $\beta \subset N$ with endpoints $P_1 \in \partial(M_1)$ and $P_2 \in \partial(M_2)$ and otherwise disjoint from M. We will always assume that β lies in an open analytic arc β_0 in N such that $\beta_0 - \beta \subset M$ and β_0 meets transversally $\partial(M_i)$ at P_i, $i = 1, 2$. See Figure 1.

A proper region $V \subset N$ is said to be an *annular extension* of $M \cup \beta$ in N if $M \cup \beta \subset V^\circ$, $V - (M^\circ \cup \beta)$ contains no connected components with compact closure that are disjoint from $\partial(V)$, and $V - (M^\circ \cup \beta)$ consists of a finite collection of compact annulus and once punctured closed discs (see Figure 2). In particular, the induced homomorphism $j_* : H_1(M \cup \beta, \mathbb{Z}) \to H_1(V, \mathbb{Z})$ is an isomorphism, where $j : M \cup \beta \to V$ is the inclusion map. If in addition the closure of $V - (M^\circ \cup \beta)$ is compact, then V is said to be a *trivial* annular extension of $M \cup \beta$.

A map $X : M \cup \beta \to \mathbb{R}^3$ is said to be *smooth* is $X|_{M_j}$, $j = 1, 2$, and $X|_{\beta_0}$ are smooth. For instance, if $Y : N \to \mathbb{R}^3$ is a smooth map then $X = Y|_{M \cup \beta}$ is smooth.

Definition 4.1 *We denote by $\mathcal{E}_N(M \cup \beta)$ the space of smooth maps $X : M \cup \beta \to \mathbb{R}^3$ such that $X_j := X|_{M_j} \in \mathcal{E}_N(M_j)$, $j = 1, 2$, and $X|_\beta$ is a smooth immersion.*



Figure 1: The surface M and the curve β when $M_1 \cap M_2 = \emptyset$ and $M_1 = M_2$.



Figure 2: An annular neighborhood V of $M \cup \beta$.

It is clear that $Y|_{M\cup\beta} \in \mathcal{E}_N(M \cup \beta)$ for all $Y \in \mathcal{E}(N)$.

Consider $X \in \mathcal{E}_N(M \cup \beta)$ and let $\sigma : \beta \to \mathbb{R}^3$ be a *smooth normal field* along β with respect to X. This simply means that σ extends as a smooth and unit field to β_0, $\sigma(\beta_0(t))$ is orthogonal to $(X|_{\beta_0})'(t)$ for any smooth parameter t on $X|_{\beta_0}$, and σ is tangent to X at points of $M \cap \beta_0$ (see Figure 3). Let s be an arclength parameter along $X|_\beta$ and label $B_s(P_i)$ as the ordered basis $\{(X|_\beta)'(s_i), \sigma(s_i)\}$ of the tangent plane of $X|_M$ at $P_i = \beta(s_i)$, $i = 1, 2$. The field σ is said to be *orientable* with respect to X if $B_s(P_1)$ and $B_s(P_2)$ are both positive or negative with respect to the orientation of N.

Definition 4.2 *We call* $\mathcal{E}_N^*(M \cup \beta)$ *as the space of marked immersions* $X_\sigma := (X, \sigma)$, *where* $X \in \mathcal{E}_N(M \cup \beta)$ *and* σ *is a orientable smooth normal field along* β *with respect to* X, *endowed with the* $\mathcal{C}^0(M \cup \beta)$*-topology of the uniform convergence of maps and normal fields on* $M \cup \beta$.

Given $X_\sigma \subset \mathcal{E}_N^*(M \cup \beta)$, let $\partial_z X_\sigma = (\hat{\phi}_j)_{j=1,2,3}$ be the complex vectorial 1-form of type $(1, 0)$ on $M \cup \beta$ given by $\partial_z X_\sigma|_M = \partial_z(X|_M)$, $\partial_z X_\sigma(\beta'(s)) := (X|_\beta)'(s) + i\sigma(s)$, where s is the arclength parameter of $X|_\beta$ for which $B_s(P_1)$ and $B_s(P_2)$ are both positive. To be more



Figure 3: A smooth normal field σ along β with respect to X.

precise, if $(U, z = x + iy)$ is a conformal chart on N such that $\beta \cap U = z^{-1}(\mathbb{R} \cap z(U))$, then $(\partial_z X_\sigma)|_{\beta\cap U} = \big[(X \circ \beta)'(s) + i\sigma(s)\big] s'(x) dz|_{\beta\cap U}$, and so $\partial_z X_\sigma \in \mathfrak{W}_0^{\hat{N}}(M \cup \beta)^3 \cap \mathfrak{W}^{\hat{N}}(M^c \cup \beta)^3$, where $M^c = M_1^c \cup M_2^c$ is the Osserman compactification of M and $\hat{N}$ is the open Riemann surface constructed by filling out the punctures corresponding to the annular ends of N with finite conformal type ($M^c \subset \hat{N}$ in a natural way). Likewise set $\hat{\eta}_1 = \hat{\phi}_1 - i\hat{\phi}_2$, $\hat{\eta}_2 = -\hat{\phi}_1 - i\hat{\phi}_2$ and $\hat{g} = \hat{\eta}_2/\hat{\phi}_3$. Note that $\hat{\phi}_j|_\beta$ is smooth in the above sense, $j = 1, 2, 3$, and the same holds for $\hat{\eta}_i|_\beta$, $i = 1, 2$. In a similar way, $\hat{g}$ is smooth on β and lies in $\mathfrak{F}^{\hat{N}}(M^c \cup \beta)$ provided that $\hat{g} \neq \infty$ on $\beta - \{P_1, P_2\}$. Notice that $\sum_{j=1}^3 \hat{\phi}_j^2 = 0$ and $\mathrm{Re}(\hat{\phi}_j)$ is an exact real 1-form on $M \cup \beta$, $j = 1, 2, 3$, hence we also have $X(P) = X(Q) + \mathrm{Re} \int_Q^P (\hat{\phi}_j)_{j=1,2,3}$, P, $Q \in M \cup \beta$. The pairs $(\hat{g}, \hat{\phi}_3)$ and $(\hat{\eta}_j|_M)_{j=1,2}$ will be called as the generalized Weierstrass data and spinorial representation of X_σ, respectively. As $X|_M \in \mathcal{E}_N(M)$, then $(\phi_j)_{j=1,2,3} := (\hat{\phi}_j|_M)_{j=1,2,3}$, $(\eta_j)_{j=1,2} = (\hat{\eta}_j|_M)_{j=1,2}$ and $g := \hat{g}|_M$ are the Weierstrass data, spinorial representation and meromorphic Gauss map of $X|_M$, respectively. Recall that all these data extend meromorphically to M^c.

The group homomorphism $p_{X_\sigma} : H_1(M \cup \beta, \mathbb{Z}) \to \mathbb{R}^3$, $\quad p_{X_\sigma}(\gamma) = \mathrm{Im} \int_\gamma \partial_z X_\sigma$, is said to be the *generalized flux map* of X_σ.

Definition 4.3 *For any $Y \in \mathcal{E}(N)$ we write σ_Y for the conormal field of Y along β, and set*

$$\mathfrak{R} : \mathcal{E}(N) \to \mathcal{E}_N^*(M \cup \beta), \quad \mathfrak{R}(Y) = (Y|_{M\cup\beta}, \sigma_Y).$$

It is easy to check that the restriction map $\mathfrak{R}$ is continuous.

For any $Y \in \mathcal{E}(N)$ we have that $p_{\mathfrak{R}(Y)} = p_Y|_{H_1(M\cup\beta)}$, where p_Y is the flux map of Y. Two marked immersions X_{σ_1}, $Y_{\sigma_2} \in \mathcal{E}_N^*(M \cup \beta)$ are said to be *homologically equivalent* on $M \cup \beta$ if $p_{X_{\sigma_1}} = p_{Y_{\sigma_2}}$.

Definition 4.4 *Given $X_\sigma \in \mathcal{E}_N^*(M \cup \beta)$, we denote by $\mathcal{E}_{X_\sigma}(N)$ the space of immersions $Y \in \mathcal{E}(N)$ for which $\mathfrak{R}(Y)$ is homologically equivalent to X_σ.*

4.1 The Algebraic Bridge Principle

Throughout this subsection, N is an open Riemann surface *with finite conformal type.* Consider M_i, $i = 1, 2$ and β as in the beginning of Section 4, and keep calling $P_i = \beta \cap M_i \in \partial(M_i)$, $i = 1, 2$, as the endpoints of β. We will also suppose that $N - (M \cup \beta)$ consists of a finite collection of pairwise disjoint once punctured discs, hence $j_* : H_1(M \cup \beta, \mathbb{Z}) \to H_1(N, \mathbb{Z})$ is an isomorphism, where $j : M \cup \beta \to N$ is the inclusion map.

Write ν for the genus of the Osserman compatification N^c of N, and put $M^c := M_1^c \cup M_2^c \subset N^c$, $M = M^c - \{E_1, \ldots, E_a\}$, $N = N^c - \{E_1, \ldots, E_{a+b}\}$ and $N_0 = N^c - \{E_{a+1}, \ldots, E_{a+b}\}$. Label c_i as the loop in $\partial(M_i)$ containing P_i, $i = 1, 2$, and $U_1, \ldots, U_b$ as the connected components (open discs) of $N^c - (M^c \cup \beta)$, where up to relabeling $E_{a+j} \in U_j$, $j = 1, \ldots, b$.

Let B_0 be a homology basis of $H_1(M^c \cup \beta, \mathbb{Z})$, hence of $H_1(N_0, \mathbb{Z})$. When $M_1 = M_2$ we choose B_0 so that there exists an unique $c_0 \in B_0$ such that $c_0 \cap \beta \neq \emptyset$. Furthermore, we can assume that $\beta \subset c_0$ and $c_0 - \beta$ is an open arc in $\mathrm{Int}(M^c)$ (see Figure 4).



Figure 4: The curve c_0 when $M = M_1 = M_2$ and β joins two points of the same connected component of $\partial(M)$.

The following density theorem is the core of our paper.

Theorem 4.1 (The Algebraic Bridge Principle) *Let X_σ be an arbitrary immersion in $\mathcal{E}_N^*(M \cup \beta)$. Then there exists a sequence $\{Y_n\}_{n \in \mathbb{N}} \subset \mathcal{E}_{X_\sigma}(N)$ such that $\{(Y_n|_{M \cup \beta}, \sigma_{Y_n})\}_{n \in \mathbb{N}} \to X_\sigma$ in the $\mathcal{C}^0(M \cup \beta)$-topology. Furthermore, if C is a positive constant and V an annular extension of $M \cup \beta$ in N, $\{Y_n\}_{n \in \mathbb{N}}$ can be chosen in such a way that $d_{Y_n}(M \cup \beta, \partial(V)) \geq C$ for all n, where d_{Y_n} is the intrinsic distance in N induced by Y_n.*

The global strategy for proving this theorem has essentially two phases. Firstly, we show that the spinorial representation of X_σ on $M \cup \beta$ can be approximated by holomorphic spinorial data on N extending meromorphically to N^c. These technical results are collected in Propositions 4.1 and 4.2 and Lemma 4.1. In a second step, we prove that the approximate meromorphic data on N^c can be slightly deformed in order to solve the period problem. This part corresponds to Lemmae 4.2 and 4.3. Finally we prove Theorem 4.1.

Let's start with the first phase of the program.

As usual, write $X_\sigma = (X, \sigma)$ and $X_i := X|_{M_i}$, $i = 1, 2$.

Proposition 4.1 *There exists a sequence $\{Y_{\sigma_j}^j\}_{j \in \mathbb{N}} \subset \mathcal{E}_N^*(M \cup \beta)$ such that $\{Y_{\sigma_j}^j\}_{j \in \mathbb{N}} \to X_\sigma$ uniformly on $M \cup \beta$, $Y_{\sigma_j}^j$ is homologically equivalent to X_σ on $M \cup \beta$ and $Y^j|_{M_i}$ is non flat, $j \in \mathbb{N}$.*

Proof: The proof is direct, we only give a brief sketch of it. Since any flat minimal surface can be approximated by non flat ones, we can find $\{Y^j\}_{j \in \mathbb{N}} \subset \mathcal{E}_N(M)$ such that $Y^j|_{M_i}$ is non flat for all j and $\{Y^j\}_{j \in \mathbb{N}} \to X|_M$ in the $\mathcal{C}^0(M)$ topology. Then, we extend smoothly Y^j to β and construct a smooth normal field $\sigma_j : \beta \to \mathbb{R}^3$ with respect to Y^j in such a way that $Y_{\sigma_j}^j$ is homologically equivalent to X_σ and $(Y^j|_\beta, \sigma_j)\}_{j \in \mathbb{N}} \to (X|_\beta, \sigma)$ in the $\mathcal{C}^0$ topology on β. □

By Proposition 4.1, we can restrict ourselves to the case that X_i is non-flat, $i = 1, 2$. Indeed, if the theorem held in the non flat case, the immersions $Y_{\sigma_j}^j$ in Proposition 4.1 would lie in the closure of $\mathfrak{R}(\mathcal{E}_{X_\sigma}(N))$ in $\mathcal{E}_N^*(M \cup \beta)$, were $\mathfrak{R}$ is the restriction map in Definition 4.3, hence the same would occur for X and the first part of the theorem would hold. The second one would be also obtained in the same process.

Label $\partial_z X_\sigma = (\hat{\phi}_j)_{j=1,2,3}$, and consider the generalized Weierstrass data $(\hat{g}, \hat{\phi}_3)$ and spinorial representation $(\hat{\eta}_j|_M)_{j=1,2}$ of X_σ. Write $d\hat{g}$ for the 1-form of type $(1,0)$ on $M^c \cup \beta$ given by $d\hat{g}|_{M^c} = d(\hat{g}|_{M^c})$ and $d\hat{g}(\beta'(s)) = (\hat{g}|_\beta)'(s)$, where s is the arclength parameter of $X|_\beta$. In other words, if $(U, z = x+iy)$ is a conformal chart in N so that $\beta \cap U = z^{-1}(\mathbb{R} \cap z(U))$, then $d\hat{g}|_{\beta\cap U} = (\hat{g}|_\beta)'(s)s'(x)dz|_{\beta\cap U}$, and so $d\hat{g} \in \mathfrak{W}^{N^c}(M^c \cup \beta)$ when $\hat{g}(P) \neq \infty$ for all $P \in \beta - \{P_1, P_2\}$. Write $(\phi_j)_{j=1,2,3} = (\hat{\phi}_j|_M)_{j=1,2,3}$, $(\eta_j)_{j=1,2} = (\hat{\eta}_j|_M)_{j=1,2}$ and $g = \hat{g}|_M$ for the Weierstrass data, the spinorial representation and the meromorphic Gauss map of $X|_M$, respectively, and call with the same name their meromorphic extensions to M^c.

Proposition 4.2 *Without loss of generality, we can assume that $\hat{g}$, $1/\hat{g}$, $(\hat{g}^2-1)$, and $d\hat{g}$ never vanish on $\partial(M) \cup \beta$ (hence the same holds for $\hat{\eta}_i$, $i = 1,2$, and $\hat{\phi}_j$, $j = 1,2,3$). In particular, $\hat{g} \in \mathfrak{F}^{N^c}(M^c \cup \beta)$ and $d\hat{g} \in \mathfrak{W}^{N^c}(M^c \cup \beta)$.*

Likewise, we can also assume that $g(E_i) \neq 0, \infty$, $i = 1, \ldots, a$, and $g^{-1}(\{0,\infty\})$ contains no branch points of g. Therefore, $m_i := Ord_{E_i}(\phi_3) = Ord_{E_i}(\eta_1) = Ord_{E_i}(\eta_2) > 1$, where $Ord_{E_i}(\cdot)$ means pole order at E_i, $i = 1, \ldots, a$.

Proof: Up to a rigid motion, we can suppose that $g(E_i) \neq 0, \infty$, $i = 1, \ldots, a$ and $g^{-1}(\{0,\infty\})$ contains no branch points of g. Recall that $X|_M$ is non flat and extends as a conformal minimal immersion beyond M in N. Therefore, we can find a sequence $M_{i,1} \supset M_{i,2} \supset \ldots$ of trivial annular extensions of M_i in N such that $M_{i,n} \subset M_{i,n-1}^\circ$ for any n, $\cap_{n\in\mathbb{N}} M_{i,n} = M_i$, X_i and g extend (with the same name) as a conformal minimal immersion and a meromorphic function to $M_{i,1}$, and g, $1/g$, (g^2-1), and dg never vanish on $\partial(M_{i,n})$ for all n, $i = 1,2$. Label $N_n = M_{1,n} \cup M_{2,n}$, and without loss of generality assume that $\beta_n := \beta - N_n^\circ$ is a Jordan arc connecting $\partial(M_{1,n})$ and $\partial(M_{2,n})$. Up to suitably deforming $X|_\beta$ and $\sigma|_\beta$, we can construct marked immersions $Z^n_{\sigma_n} \in \mathcal{E}^*_N(N_n \cup \beta_n)$ homologically equivalent to X_σ on $M \cup \beta$ such that $Z^n|_{M_{i,n}} = X_i|_{M_{i,n}}$, $i = 1,2$, $\{Z^n_{\sigma_n}|_{M\cup\beta}\}_{n\in\mathbb{N}} \to X_\sigma$ in the $\mathcal{C}^0(M \cup \beta)$-topology, and $\hat{g}_n$, $1/\hat{g}_n$, $(\hat{g}_n^2 - 1)$, and $d\hat{g}_n \neq 0$ on $\partial(N_n) \cup \beta_n$, where $\hat{g}_n$ is the generalized Gauss map of Z^n. If the theorem held for $Z^n_{\sigma_n}$, $n \in \mathbb{N}$, we would infer that $Z^n_{\sigma_n}$ lies in the closure of $\mathfrak{R}(\mathcal{E}_{X_\sigma}(N))$ in $\mathcal{E}^*_N(N_n \cup \beta_n)$, $n \in \mathbb{N}$, hence the same would occur for $X|_{M\cup\beta}$. The second part of the theorem would be also obtained in the same process. $\Box$

Let Θ_j denote the class of η_j in $\frac{\mathfrak{S}(M)}{\sim}$, and for simplicity write ξ_j for $\xi_{\Theta_j} : H_1(M,\mathbb{Z}) \to \mathbb{Z}_2$. Observe that these objects make sense even when $M_1 \cap M_2 = \emptyset$ and M is not connected. Let us show that there is a natural extension of ξ_j to $H_1(N,\mathbb{Z})$ depending on $\hat{\eta}_j$. If $M_1 \cap M_2 = \emptyset$ then $H_1(N,\mathbb{Z}) = H_1(M \cup \beta, \mathbb{Z}) = H_1(M_1,\mathbb{Z}) \oplus H_1(M_2,\mathbb{Z})$ and the extension is trivial, hence only the case $M = M_1 = M_2$ requires an explanation. Recall that all the zeroes of $\eta_j = \hat{\eta}_j|_M$ have even order and $\hat{\eta}_j$ never vanishes on $\partial(M) \cup \beta$ from Proposition 4.2, $j = 1,2$. Take any conformal chart (A, z), where $A \subset N$ is an annular neighborhood of c_0, and simply set $\xi_j(c_0) = 0$ if $\sqrt{\hat{\eta}_j(z)/dz}$ has a well defined branch along c_0 and $\xi_s(c_0) = 1$ otherwise.

The fact that $\hat{\eta}_2/\hat{\eta}_1 = \hat{g}^2$ implies $\xi_1 = \xi_2$, and in this sense $\hat{\eta}_1$ and $\hat{\eta}_2$ are spinorially equivalent on $\mathrm{Int}(M) \cup \beta$. Lemma 2.1 guarantees the existence of a unique spinor structure on N associated to ξ_1. By definition, an 1-form $\theta \in \mathfrak{S}(N)$ is said to be spinorially equivalent to $\hat{\eta}_1$ (and so to $\hat{\eta}_2$) if $\xi_\Theta = \xi_1$, where $\Theta \in \frac{\mathfrak{S}(N)}{\sim}$ is the class of θ.

Lemma 4.1 *There are $\{\eta_1^n\}_{n\in\mathbb{N}}$, $\{\eta_2^n\}_{n\in\mathbb{N}} \subset \mathfrak{S}_0(N) \cap \mathfrak{W}(N^c)$ such that:*

(i) $\{\eta_j^n|_{M^c\cup\beta}\}_{n\in\mathbb{N}} \to \hat{\eta}_j$ in the $\omega(M^c \cup \beta)$-topology, η_j^n never vanishes on $\partial(M) \cup \beta$, $(\eta_j^n|_{M^c}) = (\hat{\eta}_j)$, $(\eta_j^n|_{M^c} - \hat{\eta}_j)_0 \geq \prod_{i=1}^a E_i^{m_i}$, and $(\eta_j^n)_\infty \geq \prod_{k=a+1}^{a+b} E_k$, $j \in \{1,2\}$, for all $n \in \mathbb{N}$, $j = 1,2$.

(ii) η_1^n and η_2^n are spinorially equivalent in $\mathfrak{S}(N)$ and have no commom zeroes on N.

Proof: The following claims will be useful:

> **Claim 4.1** *Let R be a compact Riemann surface with empty boundary and genus ν. Given an open disc $U \subset R$, a point $Q \in R$ and a divisor $D_1 \in \mathfrak{Div}(R)$, there exists an integral divisor $D_2 \in \mathfrak{Div}(U)$ of degree ν and $n_0 \in \mathbb{N}$ such that $D_2^{n_0} D_1^{-1} Q^{-m}$ is the principal divisor associated to some $f \in \mathfrak{F}(R)$, where $m = n_0\nu - \mathfrak{Deg}(D_1)$.*

Proof: Since the proof is trivial when $\nu = 0$, we will assume that $\nu \geq 1$. By Jacobi's theorem, we can find an open disc $W \subset U$ such that $\varphi_Q : W_\nu \to \varphi_Q(W_\nu)$ is a diffeomorphism, where φ_Q is the Abel-Jacobi map with base point Q and W_ν is the set of divisors in R_ν with support in U. Since $J(R)$ is a compact additive Lie Group and $\varphi_Q(W_\nu) \subset J(R)$ is an open subset, for large enough $n_0 \in \mathbb{N}$ one has $n_0\varphi_E(W_\nu) = J(R)$. Therefore, there is $D_2 \in W_\nu$ such that $\varphi_Q(D_2^{n_0}) = \varphi_Q(D_1) = \varphi_Q(D_1 Q^m)$, where $m = n_0\nu - \mathfrak{Deg}(D_1)$. The lemma follows from Abel's theorem. □

> **Claim 4.2** *We can find θ_1, $\theta_2 \in \mathfrak{S}_0(N) \cap \mathfrak{W}(N^c)$ so that $|\theta_1| + |\theta_2|$ has no zeroes in N, θ_j is spinorially equivalent to $\hat{\eta}_1$ and $\hat{\eta}_2$, $\theta_j|_{M\cup\beta}$ is never-vanishing and $(\theta_j)_\infty \geq \prod_{i=1}^a E_i^{2m_i}$, $j = 1, 2$.*

Proof: Take $\theta \in \mathfrak{S}(N)$ spinorially equivalent to $\hat{\eta}_1$ and $\hat{\eta}_2$ (see Lemma 2.1).

Let k_i denote the zero order of θ at E_i ($k_i = 0$ provided that $\theta(E_i) \neq 0$), $i = 1, \ldots, a$, write $(\theta|_N) = D_0^2$, and fix two disjoint open discs V_1, $V_2 \subset N^c - (M^c \cup \beta)$. By Claim 4.1, there are $D_j \in \mathfrak{Div}(V_j)$, $n_j \in \mathbb{N}$ and $h_j \in \mathfrak{F}(N^c)$ such that $(h_j) = D_j^{n_j} E_{a+b}^{-v_j} D_0^{-1} \prod_{i=1}^a E_i^{-m_i-k_i}$, where $v_j = n_j\mathfrak{Deg}(D_j) - \mathfrak{Deg}(D_0) - \sum_{i=1}^a (m_i + k_i)$, $j = 1, 2$. It suffices to put $\theta_j = h_j^2\theta$, $j = 1, 2$. □

Let $m_{i,j} \geq 2m_i$ denote the pole order of θ_j at E_i, $i = 1, \ldots, a$, and likewise call $n_{k,j}$ as the zero order of θ_j at E_{a+k} ($n_{k,j} = 0$ provided that $\theta_j(E_{a+k}) \neq 0$), $k = 1, \ldots, b$. Set $s_j = \frac{\hat{\eta}_j}{\theta_j}$, $j = 1, 2$, and observe that: $s_j \in \mathfrak{F}_0^{N^c}(M^c \cup \beta)$. Moreover, Proposition 4.2 and Claim 4.2 give that $s_j \neq 0, \infty$ on $\partial(M) \cup \beta$, $(s_j|_{M^c}) = (\hat{\eta}_j)_0 \prod_{i=1}^a E_i^{m_{i,j}-m_i} \geq \prod_{i=1}^a E_i^{m_i}$, $j = 1, 2$, and $|s_1| + |s_2| \neq 0$ has no zeroes in $M \cup \beta$. By Claim 4.2 again, there is $t_j \in \mathfrak{F}_0^{N^c}(M^c \cup \beta)$ such that $s_j = t_j^2$, $j = 1, 2$.

Let C_1 be a collection of pairwise disjoint closed discs in $N - (M^c \cup \beta)$ containing all the zeroes of $\theta_2|_N$ (see Claim 4.2) and meeting all the bounded components of $N^c - (M^c \cup \beta)$.

Consider a continuous map $t_1^* : M^c \cup \beta \cup C_1 \to \mathbb{C}$, $t_1^*|_{M^c\cup\beta} = t_1$ and $t_1^*|_{C_1} = \delta$, where δ is any non-zero constant. By Theorem 2.1 and for each $n \in \mathbb{N}$ we can find a function $H_{n,1} \in \mathfrak{F}(N^c) \cap \mathfrak{F}_0(N_0)$ such that $|H_{n,1} - t_1^*| < 1/n$ on $M^c \cup \beta \cup C_1$ and $(H_{n,1}|_{M^c} - t_1)_0 \geq (\hat{\eta}_j)_0 \prod_{i=1}^a E_i^{m_{i,1}+m_i}$. In particular, $(H_{n,1}/t_1)|_{M^c}$ is holomorphic and $(H_{n,1}/t_1)|_{M^c\cup\beta}$ and $H_{n,1}|_{C_1}$ are never-vanishing for large enough n (without loss of generality, for all n).

Let us see that the sequence of pole orders $\{\mathrm{Ord}_{E_{a+k}} H_{n,1}\}_{n\in\mathbb{N}}$ is divergent for all $k \in \{1, \ldots, b\}$. Reason by contradiction, and assume that this sequence is bounded for some $k \in \{1, \ldots, b\}$. Since $\{H_{n,1}|_{\partial(\overline{U}_k)}\}_{n\in\mathbb{N}}$ converges uniformly to $t_1|_{\partial(\overline{U}_k)}$ and $H_{n,1} \in \mathfrak{F}_0^{N^c}(\overline{U}_k - \{E_{a+k}\}) \cap \mathfrak{F}^{N^c}(\overline{U}_k)$ for all n, up to subsequences $\{H_{n,1}|_{\overline{U}_k}\}_{n\in\mathbb{N}} \to w_0 \in \mathfrak{F}^{N^c}(\overline{U}_k) \cap \mathfrak{F}_0^{N^c}(\overline{U}_k - \{E_{a+k}\})$, where $w_0|_{\partial(\overline{U}_k)} = t_1|_{\partial(\overline{U}_k)}$. However $\{H_{n,1}|_{U_k\cap C_1}\}_{n\in\mathbb{N}} \to \delta$ and $w_0|_{\partial(\overline{U}_k)}$ is non-constant, contradicting the Identity Principle. Thus, without loss of generality we can assume that $\mathrm{Ord}_{E_k} H_{n,1} \geq (n_{k,1} + 1)/2$ for all $n \in \mathbb{N}$, $k = 1, \ldots, b$.

Call $F_{n,1} = (H_{n,1})^2$, and notice that $\{F_{n,1}\}_{n\in\mathbb{N}} \to s_1$ in the $\omega(M^c \cup \beta)$-topology, $F_{n,1}/s_1$ is never-vanishing on $M^c \cup \beta$ and holomorphic on M^c, $F_{n,1}$ never vanishes on $\partial(M) \cup \beta$, $(F_{n,1}\theta_1|_{M^c}) = (\hat{\eta}_1)$, $\big((F_{n,1} - s_1)|_{M^c}\big)_0 \geq (\hat{\eta}_1)_0 \prod_{i=1}^a E_i^{m_{i,1}+m_i}$ and $(F_{n,1})_\infty \geq \prod_{k=a+1}^{a+b} E_k^{n_{k,1}+1}$ for all n.

Setting $\eta_1^n := F_{n,1}\theta_1$, item (i) holds for $j = 1$.

Choose a collection $C_{n,2}$ of pairwise disjoint closed discs in $N - (M^c \cup \beta)$ containing all the zeroes of η_1^n in $N - M^c$ and meeting all the bounded components of $N^c - (M^c \cup \beta)$. Set $t_{n,2}^* : M^c \cup \beta \cup C_{n,2} \to \mathbb{C}$, $t_{n,2}^*|_{M^c\cup\beta} = t_2$ and $t_{n,2}^*|_{C_{n,2}} = \delta$, where δ is any non-zero constant. As above we can construct $H_{n,2} \in \mathfrak{F}(N^c) \cap \mathfrak{F}_0(N_0)$ satisfying that $|H_{n,2} - t^*_{n,2}| < 1/n$ on $M^c \cup \beta \cup C_{n,2}$, $(H_{n,2}|_{M^c} - t_2)_0 \geq (\hat{\eta}_2)_0 \prod_{i=1}^a E_i^{m_{i,2}+m_i}$, $H_{n,2}/t_2$ is holomorphic and never-vanishing on $M^c \cup \beta$, $H_{n,2}|_{C_{n,2}}$ is holomorphic never vanishing, and $\mathrm{Ord}_{E_k} H_{n,2} \geq (n_{k,2}+1)/2$ for all $n \in \mathbb{N}$, $k = 1, \ldots, b$. Set $F_{n,2} = (H_{n,2})^2$, and observe that $\{F_{n,2}\}_{n\in\mathbb{N}} \to s_2$ in the $\omega(M^c \cup \beta)$-topology on $M^c \cup \beta$, the holomorphic function $F_{n,2}/s_2$ is never-vanishing on $M^c \cup \beta$ for all n, $(F_{n,2}\theta_2|_{M^c}) = (\hat{\eta}_2)$, $\big((F_{n,2} - s_2)|_{M^c}\big)_0 \geq (\hat{\eta}_2)_0 \prod_{i=1}^a E_i^{m_{i,2}+m_i}$ and $(F_{n,2})_\infty \geq \prod_{k=a+1}^{a+b} E_k^{n_{k,2}+1}$ for all n.

Choosing $\eta_2^n := F_{n,2}\theta_2$, item (i) holds for $j = 2$.

Let us check item (ii). Obviously, η_1^n and η_2^n are spinorially equivalent in $\mathfrak{S}(N)$. Note that $F_{n,1}|_{C_1}$ never vanishes, hence $|\eta_1^n| + |\theta_2|$ has no zeroes on N from the choice of C_1, $n \in \mathbb{N}$. Likewise, the choice of $C_{n,2}$ and the fact that $F_{n,2}|_{C_{n,2}}$ never vanishes imply that $|\eta_1^n| + |\eta_2^n|$ have no zeroes on $N - (M \cup \beta)$. Moreover, η_1^n and η_2^n never vanishes on $\partial(M) \cup \beta$ and $(\eta_j^n|_{M^c}) = (\hat{\eta}_j)$, $j = 1, 2$, hence $|\eta_1^n| + |\eta_2^n|$ never vanishes on M too and we are done. $\square$

By (ii) in Lemma 4.1, $\phi_3^n := \sqrt{\eta_1^n \eta_2^n}$ is well defined and lies in $\mathfrak{W}(N^c) \cap \mathfrak{W}_0(N)$. Up to a suitable choice of the branch, we will assume that $\phi_3^n|_M - \phi_3$ extends holomorphically to M^c and $\{\phi_3^n|_{M^c\cup\beta} - \hat{\phi}_3\}_{n\in\mathbb{N}} \to 0$ on $M^c \cup \beta$ as well.

At this point we can start with the second phase of the program.

Set $\mathcal{L} = \{f \in \mathfrak{F}(N^c) \cap \mathfrak{F}_0(N_0) \ : \ (f)_0 \geq \prod_{j=1}^a E_j^{m_i}\}$. By Riemann-Roch theorem, $\mathcal{L}$ is a linear subspace of $\mathfrak{F}(N^c) \cap \mathfrak{F}_0(N_0)$ with infinite dimension (and finite codimension).

Consider the complex normed space $(\mathfrak{F}_0^{N^c}(M^c \cup \beta), \|\cdot\|)$, where $\|h\| = \mathrm{Max}\{|h(P)| \ : \ P \in M^c \cup \beta\}$ is the norm of the maximum inducing the $\omega(M^c \cup \beta)$-topology on $M^c \cup \beta$. Up to the restriction to $M^c \cup \beta$, we can consider $\mathcal{L} \subset \mathfrak{F}_0^{N^c}(M^c \cup \beta)$.

For each $f \in \mathfrak{F}_0^{N^c}(M^c \cup \beta)$ set $\hat{\eta}_j(f) = (1 + jf)^2 \hat{\eta}_j$, $\quad j = 1, 2$, likewise define $\hat{\phi}_k(f)$ and notice that $\hat{\phi}_k(f) - \hat{\phi}_k \in \mathfrak{F}_0^{N^c}(M^c \cup \beta)$, $k = 1, 2, 3$. Call $\varsigma_0 = 3(2\nu + b - 1)$ and consider the Fréchet differentiable map:

$$\mathcal{P} : \mathfrak{F}_0^{N^c}(M^c \cup \beta) \to \mathbb{C}^{\varsigma_0}, \ \mathcal{P}(f) = [(\int_d \hat{\phi}_j(f) - \hat{\phi}_j)_{d\in B_0}]_{j=1,2,3}.$$

It is clear that $\mathcal{P}(\mathbf{0}) = [(0)_{d\in B_0}]_{j=1,2,3}$, where $\mathbf{0}$ is the constant zero function.

Lemma 4.2 *If $d\mathcal{P}_0 : \mathfrak{F}_0^{N^c}(M^c \cup \beta) \to \mathbb{C}^{\varsigma_0}$ denotes the complex Fréchet derivative of $\mathcal{P}$ at $\mathbf{0}$ then $d\mathcal{P}_0(\mathcal{L}) = \mathbb{C}^{\varsigma_0}$.*

Proof: Reason by contradiction, and assume that $d\mathcal{P}_0(\mathcal{L})$ lies in a hyperplane $U = \{((x_d^j)_{d\in B_0})_{j=1,2,3} \in \mathbb{C}^{\varsigma_0} \ : \ \sum_{j=1}^3 (\sum_{d\in B_0} \lambda_d^j x_d^j) = 0\}$, where $\sum_{j=1}^3 (\sum_{d\in B_0} |\lambda_d^j|) \neq 0$.

Therefore $d\mathcal{P}_0(f) = \frac{d\mathcal{P}(tf)}{dt}|_{t=0} \in U$, for any $f \in \mathcal{L}$, that is to say

$$\int_{\Gamma_1} f\hat{\eta}_1 + \int_{\Gamma_2} f\hat{\eta}_2 + \int_{\Gamma_3} f\hat{\phi}_3 = 0, \text{ for all } f \in \mathcal{L}. \tag{3}$$

where $\Gamma_j \in H_1(M^c \cup \beta, \mathbb{C})$, $j = 1, 2, 3$, are the cycles with complex coefficients given by:

$$\Gamma_1 = \sum_{d\in B_0} (\lambda_d^1 + i\lambda_d^2)\, d, \ \Gamma_2 = 2\sum_{d\in B_0} (-\lambda_d^1 + i\lambda_d^2)\, d, \ \Gamma_3 = 3\sum_{d\in B_0} \lambda_d^3\, d.$$

In the remaining of the proof, we consider ϕ_k, η_j as meromorphic 1-forms in $\mathfrak{W}^{N^c}(M^c)$ and g as a meromorphic function in $\mathfrak{F}^{N^c}(M^c)$. Furthermore, we only deal with divisors of meromorphic functions and 1-forms *relative to* M^c. It is crucial to take into account Proposition 4.2.

Set $\mathcal{L}_0=\{f\in\mathcal{L}\ :\ (f)_0\geq(\phi_3)_0^2=(g)_0^2(g)_\infty^2\}$ and note that $\mathcal{L}_0$ is a linear subspace of $\mathcal{L}$ of of infinite dimension from Riemann-Roch theorem. Since $m_i=\mathrm{Ord}_{E_i}(\phi_3)$ and $g(E_i)\neq 0,\infty$ for all $i=1,\ldots,a$, then $df/\hat{\phi}_3\in\mathfrak{F}_0^{N^c}(M^c\cup\beta)$ and $(df/\phi_3)\geq\prod_{i=1}^a E_i^{m_i}$ for all $f\in\mathcal{L}_0$. By Theorem 2.1 there is $\{f_n\}_{n\in\mathbb{N}}\subset\mathcal{L}$ converging to $df/\hat{\phi}_3$ in the $\omega(M^c\cup\beta)$-topology. Applying equation (3) to f_n and taking the limit as n goes to $+\infty$, we infer that $\int_{\Gamma_1}\frac{df}{\hat{g}}+\int_{\Gamma_2}\hat{g}df=0$ for any $f\in\mathcal{L}_0$. Integrating by parts,

$$\int_{\Gamma_1}\frac{fd\hat{g}}{\hat{g}^2}-\int_{\Gamma_2}fd\hat{g}=0,\quad\text{for all } f\in\mathcal{L}_0. \tag{4}$$

Denote by $\mathcal{L}_1=\{f\in\mathcal{L}\ :\ (f)_0\geq(g^2-1)_0^2(dg)_0^2\prod_{i=1}^a E_i^{2m_i}\}$, and as above observe that $\mathcal{L}_1$ is a linear subspace of $\mathcal{L}$ of infinite dimension. For any $f\in\mathcal{L}_1$, the function $h_f:=\frac{\hat{g}^2df}{(\hat{g}^2-1)d\hat{g}}$ lies in $\mathfrak{F}_0^{N^c}(M^c\cup\beta)$ and satisfies that $(h_f)\geq(\phi_3)_0^2\prod_{i=1}^a E_i^{m_i}$ (take into account Proposition 4.2). By Theorem 2.1, h_f lies in the closure of $\mathcal{L}_0$ in $(\mathfrak{F}_0^{N^c}(M^c\cup\beta),\|\cdot\|)$, hence equation (4) can be formally applied to h_f to obtain that $\int_{\Gamma_1-\Gamma_2}\frac{df}{\hat{g}^2-1}=0$, for any $f\in\mathcal{L}_1$. Integrating by parts,

$$\int_{\Gamma_1-\Gamma_2}\frac{fd\hat{g}}{(\hat{g}^2-1)^2}=0,\quad\text{for all } f\in\mathcal{L}_1. \tag{5}$$

Let us show that $\Gamma_1=\Gamma_2$. Indeed, it is well known (see [4]) that there exist $2\nu+b-1$ cohomologically independent meromorphic 1-forms in $\mathfrak{W}(N^c)\cap\mathfrak{W}_0(N_0)$ generating the first holomorphic De Rham cohomology group $\mathcal{H}^1_{\mathrm{hol}}(N_0)$ of N_0. Thus, the map $H^1_{\mathrm{hol}}(N_0)\longrightarrow\mathbb{C}^{2\nu+b-1}$, $[\tau]\mapsto\left(\int_d\tau\right)_{d\in B_0}$, is a linear isomorphism. Assume that $\Gamma_1\neq\Gamma_2$ and take $\tau\in\mathfrak{W}(N^c)\cap\mathfrak{W}_0(N_0)$ such that $\int_{\Gamma_1-\Gamma_2}\tau\neq 0$. Since $\mathcal{V}:=\mathfrak{F}(N^c)\cap\mathfrak{F}_0(N_0)$ has infinite dimension, we can find $F\in\mathcal{V}$ such that $(\tau+dF)_0\geq(d\hat{g})_0^3(\hat{g})_\infty^2\prod_{i=1}^a E_i^{2m_i}$. Set $h:=\frac{(\tau+dF)(\hat{g}^2-1)^2}{d\hat{g}}\in\mathfrak{F}_0^{N^c}(M^c\cup\beta)$ and note that $(h)\geq(g^2-1)_0^2(dg)_0^2\prod_{i=1}^a E_i^{2m_i}$. By Theorem 2.1, h lies in the closure of $\mathcal{L}_1$ in $(\mathfrak{F}_0^{N^c}(M^c\cup\beta),\|\cdot\|)$ and equation (5) gives that $\int_{\Gamma_1-\Gamma_2}\tau+dF=\int_{\Gamma_1-\Gamma_2}\tau=0$, a contradiction.

Turning back to equation (4) and using that $\Gamma_1=\Gamma_2$, one has

$$\int_{\Gamma_1}f(\frac{1}{\hat{g}^2}-1)d\hat{g}=0,\quad\text{for all } f\in\mathcal{L}_0. \tag{6}$$

Let us see now that $\Gamma_1=0$. Reason by contradiction and suppose that $\Gamma_1\neq 0$. As above, take $\tau\in\mathfrak{W}(N^c)\cap\mathfrak{W}_0(N_0)$ and $H\in\mathcal{V}$ such that $\int_{\Gamma_1}\tau\neq 0$ and $(\tau+dH)_0\geq(d\hat{g})_0(\hat{g}^2-1)_0\prod_{i=1}^a E_i^{m_i}$. Set $t:=\frac{(\tau+dH)\hat{g}^2}{(\hat{g}^2-1)d\hat{g}}\in\mathfrak{F}_0^{N^c}(M^c\cup\beta)$ and observe that $(t)\geq(\phi_3)_0^2$. By Theorem 2.1, t lies in the closure of $\mathcal{L}_0$ in $(\mathfrak{F}_0^{N^c}(M^c\cup\beta),\|\cdot\|)$, hence from equation (6) we get that $\int_{\Gamma_1}(\tau+dH)=\int_{\Gamma_1}\tau=0$, a contradiction.

Finally, equation (3) and the fact that $\Gamma_1=\Gamma_2=0$ give that

$$\int_{\Gamma_3}f\hat{\phi}_3=0\quad\text{for all } f\in\mathcal{L}. \tag{7}$$

Since $\sum_{j=1}^3(\sum_{d\in B_0}(\lambda_d^j)^2)\neq 0$ and $\Gamma_1=\Gamma_2=0$, necessarily $\Gamma_3\neq 0$. Take $\tau\in\mathfrak{W}(N^c)\cap\mathfrak{W}_0(N_0)$ and $G\in\mathcal{V}$ such that $\int_{\Gamma_3}\tau\neq 0$ and $(\tau+dG)_0\geq(\phi_3)_0$. The function $v:=\frac{(\tau+dG)}{\hat{\phi}_3}$ lies in $\mathfrak{F}_0^{N^c}(M^c\cup\beta)$ and satisfies that $(v)\geq\prod_{i=1}^a E_i^{m_i}$. By Theorem 2.1, v lies in the closure of $\mathcal{L}$ in $\mathfrak{F}_0^{N^c}(M^c\cup\beta)$ and equation (7) can be formally applied to v. We get that $\int_{\Gamma_3}\tau+dF=0$, absurd. This contradiction proves the lemma. $\square$

For each $f\in\mathcal{L}$ and $n\in\mathbb{N}$ set $\eta_j^n(f)=(1+jf)^2\eta_j^n$, $j=1,2$, and likewise define $\phi_j^n(f)$, $j=1,2,3$. It is clear that $\eta_j^n(f)-\eta_j^n$, $\phi_k^n(f)-\phi_k^n\in\mathfrak{W}_0(N_0)$, hence

$$\phi_k^n(f)-\hat{\phi}_k\in\mathfrak{W}_0^{N^c}(M^c\cup\beta)\text{ for all } f\in\mathcal{L},\ n\in\mathbb{N}\text{ and } k\in\{1,2,3\}. \tag{8}$$

Following Lemma 4.2, let $\mathcal{U} \subset \mathcal{L}$ be a ς_0-dimensional complex linear subspace such that $d\mathcal{P}_0(\mathcal{U}) = \mathbb{C}^{\varsigma_0}$, and fix a basis $\{f_d^j \ : \ d \in B_0, \ j \in \{1,2,3\}\}$ of $\mathcal{U}$. For the sake of simplicity we write $\mathbf{f}_0 = [(f_d^j)_{d\in B_0}]_{j=1,2,3} \in \mathcal{L}^{\varsigma_0}$.

For any $\mathbf{x} = [(x_d^j)_{d\in B_0}]_{j=1,2,3} \in \mathbb{C}^{\varsigma_0}$ and $\mathbf{h} = [(h_d^j)_{d\in B_0}]_{j=1,2,3} \in \mathcal{L}^{\varsigma_0}$, we define $\mathbf{x}\cdot\mathbf{h} = \sum_{j=1}^3[\sum_{d\in B_0} x_d^j h_d^j]$. For each $n \in \mathbb{N}\cup\{0\}$ and $\mathbf{h} \in \mathcal{L}^{\varsigma_0}$, write $\mathcal{Q}_{n,\mathbf{h}} : \mathbb{C}^{\varsigma_0} \to \mathbb{C}^{\varsigma_0}$ for the vectorial degree two complex polynomial function given by

$$\mathcal{Q}_{n,\mathbf{h}}(\mathbf{x}) = [\Big(\int_d \phi_j^n(\mathbf{x}\cdot\mathbf{h}) - \hat{\phi}_j\Big)_{d\in B_0}]_{j=1,2,3},$$

where we have made the convention $\phi_j^0 = \hat{\phi}_j$ for all j. Equation (8) gives that $\mathcal{Q}_{n,\mathbf{h}}$ is well defined for all n and h.

By Lemma 4.2 the holomorphic map $\mathcal{Q}_{0,\mathbf{f}_0}$ non-zero Jacobian at the origin, hence we can find a closed Euclidean ball $K_0 \subset \mathbb{C}^{\varsigma_0}$ centered at the origin such that $\mathcal{Q}_{0,\mathbf{f}_0}|_{K_0} : K_0 \to \mathcal{Q}_{0,\mathbf{f}_0}(K_0)$ is a biholomorphism. Moreover, since $\mathcal{Q}_{0,\mathbf{f}_0}(\mathbf{0}) = \mathbf{0}$ then $\mathcal{Q}_{0,\mathbf{f}_0}(K_0)$ contains the origin as interior point. Up to choosing K_0 with small enough radius, we can also assume that

$$\|\mathbf{x}\cdot\mathbf{f}_0\| < 1 \text{ on } M^c\cup\beta \text{ for all } \mathbf{x}\in K_0. \tag{9}$$

Lemma 4.3 *We can find* $\{\mathbf{f}_n\}_{n\in\mathbb{N}} \subset \mathcal{L}^{\varsigma_0}$ *such that:*

(i) $\{\mathbf{f}_n|_{M^c\cup\beta}\}_{n\in\mathbb{N}} \to \mathbf{f}_0|_{M^c\cup\beta}$ *in the* $\omega(M^c\cup\beta)^{\varsigma_0}$*-topology.*

(ii) $\eta_j^n(\mathbf{x}\cdot\mathbf{f}_n)$ *has poles at* E_k *for all* $k\in\{1,\ldots,a+b\}$, $\mathbf{x}\in\mathbb{C}^{\varsigma_0}$ *and* $n\in\mathbb{N}$, $j=1,2$,

(iii) $\sum_{j=1}^2|\eta_j^n(\mathbf{x}\cdot\mathbf{f}_n)|$ *never vanishes on* N *for all* $n\in\mathbb{N}$ *and* $\mathbf{x}\in K_0$.

As a consequence $\{\eta_j^n(\mathbf{x}\cdot\mathbf{f}_n)|_{M\cup\beta}\}_{n\in\mathbb{N}} \to \hat{\eta}_j(\mathbf{x}\cdot\mathbf{f}_0)$ *in the* $\omega(M^c\cup\beta)^{\varsigma_0}$*-topology and* $\{\mathcal{Q}_{n,\mathbf{f}_n}\}_{n\in\mathbb{N}} \to \mathcal{Q}_{0,\mathbf{f}_0}$ *uniformly on* $\mathbf{x}\in K_0$.

Proof: By definition, it is clear that $\eta_j^n(f)$ has poles at E_k for all $k\in\{1,\ldots,a\}$ and $f\in\mathcal{L}$. By Lemma 4.1, $\{\eta_j^n|_{M^c\cup\beta} - \hat{\eta}_j\}_{n\in\mathbb{N}} \to 0$ on $M^c\cup\beta$ and $(\eta_j^n|_{M^c\cup\beta}) = (\hat{\eta}_j)$ for all n. Let C_n be a finite collection of closed discs in $N_0 - (M^c\cup\beta)$ containing all the zeroes of η_1^n and η_2^n in $N_0-(M^c\cup\beta)$ and meeting all the bounded components of $N^c - (M^c\cup\beta)$.

For each $d\in B_0$, $n\in\mathbb{N}$, and $j\in\{1,2,3\}$, set $\hat{f}_d^{j,n} : M^c\cup\beta\cup C_n \to \mathbb{C}$, $\hat{f}_d^{j,n}|_{M^c\cup\beta} = f_d^j$, $\hat{f}_d^{j,n}|_{C_n} = 0$.

By Theorem 2.1 and similar arguments to those used in the proof of Lemma 4.1, we can find a sequence $\{f_c^{j,n}(m)\}_{m\in\mathbb{N}}$ in $\mathcal{L}$ satisfying that $\{f_c^{j,n}(m)|_{M^c\cup\beta\cup C_n}\}_{m\in\mathbb{N}} \to \hat{f}_d^{j,n}$ in the $\omega(M^c\cup\beta\cup C_n)$-topology and the sequence of pole orders $\{\mathrm{Ord}_{E_{a+k}}\big(f_c^{j,n}(m)\big)\}_{m\in\mathbb{N}}$ is divergent for all $k\in\{1,\ldots,b\}$. Up to subsequences, we can assume that:

$$\mathrm{Ord}_{E_{a+k}}\big(f_{c_1}^{j_1,n}(m)\big) \neq \mathrm{Ord}_{E_{a+k}}\big(f_{c_2}^{j_2,n}(m)\big) \text{ provided that } (c_1,j_1)\neq(c_2,j_2). \tag{10}$$

Set $\mathbf{f}_n(m) = [(f_d^{j,n}(m))_{d\in B_0}]_{j=1,2,3}$, $m\in\mathbb{N}$, and take a divergent sequence $\{m_n\}_{n\in\mathbb{N}} \subset \mathbb{N}$ such that $\{\mathbf{f}_n(m_n)|_{M^c\cup\beta}\}_{n\in\mathbb{N}} \to \mathbf{f}_0|_{M^c\cup\beta}$ in the $\omega(|_{M^c\cup\beta})$-topology and $\big\{\max_{C_n}|f_d^{j,n}(m_n)|\big\}_{n\to\mathbb{N}} \to 0$ for all $d\in B_0$ and $j\in\{1,2,3\}$.

Set $\mathbf{f}_n = \mathbf{f}_n(m_n)$ for all $n\in\mathbb{N}$, and let us show that $\{\mathbf{f}_n\}_{n\in\mathbb{N}}$ solves the claim.

Items (i) is obvious, and item (ii) follows from equation (10) and the facts that η_j^n has poles at E_k for all $k\in\{a+1,\ldots,a+b\}$ and $\eta_j^n(f)$ has poles at E_k for all $k\in\{1,\ldots,a\}$ and $f\in\mathcal{L}$.

Let us check (iii). Up to removing finitely many terms of $\{\mathbf{f}_n\}_{n\in\mathbb{N}}$, we can assume that:

(a) $\|\mathbf{x}\cdot\mathbf{f}_n\| < 1$ on $M^c\cup\beta$ for all $\mathbf{x}\in K_0$ (see equation (9)), and so $1+\mathbf{x}\cdot\mathbf{f}_n$ and $2+\mathbf{x}\cdot\mathbf{f}_n$ never vanish on $M^c\cup\beta$ for all $\mathbf{x}\in K_0$,

(b) $1+\mathbf{x}\cdot\mathbf{f}_n$ and $2+\mathbf{x}\cdot\mathbf{f}_n$ never vanish on C_n for all n and $\mathbf{x}\in K_0$.

Since $|\eta_1^n|+|\eta_2^n|$ never vanishes on N, (a) and (b) give that $\sum_{j=1}^2|\eta_j^n(\mathbf{x}\cdot\mathbf{f}_n)|$ never vanishes on $M\cup\beta\cup C_n$ for all n. Taking into account that $|1+f|+|1+2f|$ never vanish on N for all $n\in\mathbb{N}$ and $f\in\mathcal{L}$ and the choice of C_n, $\sum_{j=1}^2|\eta_j^n(\mathbf{x}\cdot\mathbf{f}_n)|$ never vanishes on $N-(M\cup\beta\cup C_n)$ and we are done. $\square$

Proof of the Algebraic Bridge Principle: For the sake of simplicity, write $\mathcal{P}_n=\mathcal{Q}_{n,\mathbf{f}_n}$, $n\in\mathbb{N}\cup\{0\}$. Since the coefficients of the vectorial polynomial functions $\{\mathcal{P}_n\}_{n\in\mathbb{N}}$ converge to the ones of $\mathcal{P}_0$, $\mathcal{P}_n|_{K_0}:K_0\to\mathcal{P}_n(K_0)$ is a biholomorphism and $\mathcal{P}_n(K_0)$ contains the origin as interior point provided that n is large enough (up to removing finitely many terms, for all n).

Let $\mathbf{y}_n\in K_0$ denote the unique point satisfying $\mathcal{P}_n(\mathbf{y}_n)=\mathbf{0}$, and notice that $\lim_{n\to\infty}\mathbf{y}_n=\mathbf{0}$. Set $\rho_j^n:=\eta_j^n(\mathbf{y}_n\cdot\mathbf{f}_n)$, $j=1,2$, $\psi_k^n:=\phi_k^n(\mathbf{y}_n\cdot\mathbf{f}_n)$, $k=1,2,3$, and define

$$Y_n:N\to\mathbb{R}^3,\quad Y_n(P)=X(P_1)+\operatorname{Re}\int_{P_1}^{P}(\psi_k^n)_{k=1,2,3},\ n\in\mathbb{N},$$

where $P_1=\beta\cap M_1$. By (8) and the choice of $\mathbf{y}_n$, $\psi_k^n-\hat{\phi}_k$ is an exact 1-form in $\mathfrak{W}_0^{N^c}(M^c\cup\beta)$ and Y_n is well defined. Moreover, Lemma 4.3 and Osserman's theorem imply that $Y_n\in\mathcal{E}_{X_\sigma}(N)$. As $\rho_j^n|_{M^c\cup\beta}=(1+j(\mathbf{y}_n\cdot\mathbf{f}_n|_{M^c\cup\beta}))^2(\eta_j^n|_{M^c\cup\beta}-\hat{\eta}_j)+(1+j(\mathbf{y}_n\cdot\mathbf{f}_n)|_{M^c\cup\beta})^2\hat{\eta}_j$, then Lemma 4.1, Lemma 4.3 and the fact that $\{\mathbf{y}_n\}_{n\in\mathbb{N}}\to 0$ give that $\{\rho_j^n-\hat{\eta}_j\}_{n\to\mathbb{N}}\to 0$ in the $\omega(M^c\cup\beta)$-topology. Therefore $\{(Y_n|_{M\cup\beta},\sigma_{Y_n})\}_{n\in\mathbb{N}}\to X_\sigma$ in the $\mathcal{C}^0(M\cup\beta)$-topology, proving the first part of the Theorem 4.1.

For the second one, let V be an annular extension of of $M\cup\beta$ in N and take $C>0$. Since any annular extension of $M\cup\beta$ contains a trivial one, we can suppose without loss of generality that V is a trivial annular extension of $M\cup\beta$. Let L_n be a Jorge-Xavier type labyrinth in $V^\circ-(M\cup\beta)$ adapted to C and ψ_3^n, that is to say, a finite collection of pairwise disjoint closed discs in $V-(M\cup\beta)$ such that $\int_\gamma|\psi_3^n|>C$ for any compact arc $\gamma\subset V-L_n$ connecting $\partial(M)\cup\beta$ and $\partial(V)$ (see [8] or [15]). Consider another Jorge-Xavier type labyrinth L_n' obtained as a small trivial annular extension of L_n in $V^\circ-(M\cup\beta)$. By Theorem 2.1, there is $\{h_{n,m}\}_{m\in\mathbb{N}}\subset\mathfrak{F}(N_0)\cap\mathfrak{F}_0^{N^c}(V^c)$ such that $|h_{m,n}|<1/m$ on $M^c\cup\beta$, $|h_{m,n}-m|<1/m$ on L_n' and $(h_{n,m})_0\geq\prod_{j=1}^a E_j^{m_i}$, $m\in\mathbb{N}$.

Set $\varphi_1^{n,m}=e^{-h_{n,m}}\eta_1^n$, $\varphi_2^{n,m}=e^{h_{n,m}}\eta_2^n$, and call $\tau_1^{n,m}=1/2(\varphi_1^{n,m}-\varphi_2^{n,m})$, $\tau_2^{n,m}=i/2(\varphi_1^{n,m}+\varphi_2^{n,m})$ and $\tau_3^{n,m}=\psi_3^n$. For any $f\in\mathcal{L}$, label $\varphi_1^{n,m}(f)=e^{-h_{n,m}}\eta_1^n(f)$, $\varphi_2^{n,m}(f)=e^{h_{n,m}}\eta_2^n(f)$, and call $\tau_1^{n,m}(f)=1/2(\varphi_1^{n,m}(f)-\varphi_2^{n,m}(f))$, $\tau_2^{n,m}=i/2(\varphi_1^{n,m}(f)+\varphi_2^{n,m}(f))$ and $\tau_3^{n,m}(f)=\psi_3^n(f)$. Define $\mathcal{P}_{n,m}:\mathbb{C}^{\varsigma_0}\to\mathbb{C}^{\varsigma_0}$, $\mathcal{P}_{n,m}(\mathbf{x})=\big[(\int_d\tau_j^{n,m}(\mathbf{x}\cdot\mathbf{f}_n)-\hat{\phi}_j)_{d\in B_0}\big]_{j=1,2,3}$. Since $\{\mathcal{P}_{n,m}\}_{m\in\mathbb{N}}\to\mathcal{P}_n$ uniformly on compact subsets of $\mathbb{C}^{\varsigma_0}$, $\mathcal{P}_{n,m}|_{K_0}:K_0\to\mathcal{P}_{n,m}(K_0)$ is a biholomorphism and $0\in\mathcal{P}_{n,m}(K_0)-\partial\big(\mathcal{P}_{n,m}(K_0)\big)$ as well, m is large enough (without loss of generality for all m). Furthermore, if $\mathbf{y}_{n,m}\in K_0$ denotes the unique point satisfying $\mathcal{P}_{n,m}(\mathbf{y}_{n,m})=\mathbf{0}$, then $\lim_{m\to\infty}\mathbf{y}_{n,m}=\mathbf{y}_n$. Call $\rho_1^{n,m}=e^{-h_{n,m}}\eta_1^n(\mathbf{y}_{n,m}\cdot\mathbf{f}_n)$, $\rho_2^{n,m}=e^{-h_{n,m}}\eta_2^n(\mathbf{y}_{n,m}\cdot\mathbf{f}_n)$, $\psi_1^{n,m}=1/2(\rho_1^{n,m}-\rho_2^{n,m})$, $\psi_2^{n,m}=i/2(\rho_1^{n,m}+\rho_2^{n,m})$, $\psi_3^{n,m}=\psi_3^n(\mathbf{y}_{n,m}\cdot\mathbf{f}_n)$, and set

$$Y_{n,m}:N\to\mathbb{R}^3,\quad Y_{n,m}(P)=X(T_1)+\operatorname{Re}\int_{T_1}^{P}(\psi_k^{n,m})_{k=1,2,3}.$$

Note that $Y_{n,m}$ is well defined, $(Y_{n,m}|_{M\cup\beta},\sigma_{Y_{n,m}})\in\mathcal{E}_N^*(M\cup\beta)$ and $(Y_{n,m}|_{M\cup\beta},\sigma_{Y_{n,m}})$ and X_σ are homologically equivalent on $M\cup\beta$, $m\in\mathbb{N}$. Moreover, $\{(Y_{n,m}|_{M\cup\beta},\sigma_{Y_{n,m}})\}_{m\in\mathbb{N}}\to(Y_n|_{M\cup\beta},\sigma_{Y_n})$ in the $\mathcal{C}^0(M\cup\beta)$-topology for all n.

On the other hand and by Jorge-Xavier arguments, the facts that $\{e^{h_{m,n}}\}_{m\in\mathbb{N}}\to\infty$ uniformly on L_n' and $\{(Y_n|_{M\cup\beta},\sigma_{Y_n})\}_{n\in\mathbb{N}}\to X_\sigma$ in the $\mathcal{C}^0(M\cup\beta)$-topology imply the existence of a large enough $m_n\in\mathbb{N}$ such that $d_{Y_{n,m_n}}(M\cup\beta,\partial(V))>C$ for each $n\in\mathbb{N}$. Furthermore, the sequence

$\{m_n\}_{n\in\mathbb{N}}$ can be chosen so that $\{(H_n|_{M\cup\beta}, \sigma_{H_n})\}_{n\in\mathbb{N}} \to X_\sigma$ in the $\mathcal{C}^0(M\cup\beta)$-topology, where $H_n = Y_{n,m_n}$, $n\in\mathbb{N}$.

Unfortunately, H_n is not necessarily of FTC, and we have to work a little more. Fix a closed disc $D_0 \subset N - V$ and an analytic arc β' in $N - (V^\circ \cup D_0^\circ)$ joining $\partial(V)$ and $\partial(D_0)$, and note that $N - (V \cup D_0 \cup \beta')$ is a finite collection of pairwise disjoint once punctured open conformal discs. Set $X_n' = H_n|_{V\cup D_0\cup\beta'}$ and write σ_n' for the conormal field with respect to to X_n' along β'. By the first part of the proof, there exists $\{Z_{n,j}\}_{j\in\mathbb{N}} \subset \mathcal{E}_{(X_n')_{\sigma_n'}}(N)$ such that $\{(Z_{n,j}, \sigma_{Z_{n,j}})\}_{j\in\mathbb{N}} \to (X_n', \sigma_n')$ in the $\mathcal{C}^0(V\cup D_0\cup\beta')$-topology. Without loss of generality, we can also suppose that $d_{Z_{n,j}}(M\cup\beta, \partial(V)) > C$ for all j. Since $\{(Z_{n,j}|_{M\cup\beta}, \sigma_{Z_{n,j}})\}_{j\in\mathbb{N}} \to (H_n|_{M\cup\beta}, \sigma_{H_n})$ for all n, and $\{(H_n|_{M\cup\beta}, \sigma_{H_n})\}_{n\in\mathbb{N}} \to X_\sigma$, a standard diagonal process provides a sequence $\{Z_n\}_{n\in\mathbb{N}} \subset \mathcal{E}_{X_\sigma}(N)$ such that $\{(Z_n|_{M\cup\beta}, \sigma_{Z_n})\}_{n\in\mathbb{N}} \to X_\sigma$ in the $\mathcal{C}^0(M\cup\beta)$-topology and $d_{Z_n}(M\cup\beta, \partial(V)) \geq C$ for all n, concluding the proof. $\Box$

4.2 Main results

In this subsection we approach the problem of approximation by minimal surfaces with arbitrary conformal structure and flux map. We start with the following basic consequence of Theorem 4.1.

Corollary 4.1 (Basic Approximation) *Let N be an open Riemann surface of finite conformal type, and let M be a proper (possibly non compact) region in N with compact boundary such that $N - M$ consists of a finite collection of pairwise disjoint once punctured conformal discs.*

Then for each $X \in \mathcal{E}_N(M)$, constant $C > 0$, and annular extension V of M in N, there exists $\{Y_n\}_{n\in\mathbb{N}} \in \mathcal{E}(N)$ such that $\{Y_n|_M\}_{n\in\mathbb{N}} \to X$ in the $\mathcal{C}^0(M)$-topology, $d_{Y_n}(M, \partial(V)) \geq C$ for all n, and $p_{Y_n}|_{H_1(M,\mathbb{Z})} = p_X$.

Proof: Fix a closed disc $D_0 \subset V - M$ and an analytic arc β in $N - (V^\circ \cup D_0^\circ)$ joining $\partial(V)$ and $\partial(D_0)$, and note that $N - (V \cup D_0 \cup \beta)$ is a finite collection of pairwise disjoint once punctured open conformal discs. Take any $X_0 \in \mathcal{E}_N(D_0)$ and consider an immersion $Y \in \mathcal{E}_N(M \cup D_0 \cup \beta)$ so that $Y|_M = X$, $Y|_{V_0} = X_0$. Take also an orientable normal field σ with respect to Y along β. Applying Theorem 4.1 to Y_σ we are done. $\Box$

Given an open surface N a subset $S \subset N$, and a continuous map $X = (X_1, X_2, X_3) : S \to \mathbb{R}^3$, we write $\|X\|_0 = \sup_S(\sum_{j=1}^3 X_j^2)^{1/2}$.

We are going to obtain a wide wide generalization of this corollary (see the General Approximation Theorem below). We first need the following lemma:

Lemma 4.4 *Let N be an open Riemann surface, and let M_1 and M_2 be two proper regions in N of finite conformal type. Assume that M_2 is an extension of M_1 (possibly $M_2 = N$), and consider an immersion $X \in \mathcal{E}_N(M_1)$.*

Then, for any $C > 0$, $\epsilon > 0$, and group homomorphism extension $q : H_1(M_2, \mathbb{Z}) \to \mathbb{R}$ of p_X, there exists $Z \in \mathcal{E}_N(M_2)$ such that $\|Z|_{M_1} - X\|_0 \leq \epsilon$, $d_Z(M_1, \partial(M_2)) \geq C$ and $p_Z = q$.

Proof: Label ν_0, a_0 and b_0 as the genus, number of annular ends and number of boundary components of $K := M_2 - M_1^\circ$. We reason by induction on the Euler characteristic $\chi(K) = 2 - 2\nu_0 - a_0 - b_0 \leq 0$ of K (which makes sense even if K is not connected).

If $\chi(K) = 0$ then M_2 is an annular extension of M_1. Take a trivial annular extension M_2^* of M_2 in N, and let R be a conformal compactification of M_2^*, hence of M_1. Let $E \subset R - M_2^*$ be a finite subset meeting each connected component of $R - M_2^*$ in a unique point, and observe that $X \in \mathcal{E}_{R-E}(M_1)$. By Corollary 4.1, there is $\hat{X} \in \mathcal{E}(R - E)$ satisfying that $\|\hat{X}|_{M_1} - X\|_0 \leq \epsilon$ and $d_{\hat{X}}(M_1, \partial(M_2)) \geq C$. Set $Z = \hat{X}|_{M_2}$ and notice that $Z \in \mathcal{E}_N(M_2)$. Taking into account that the homomorphism $j_* : H_1(M_1, \mathbb{Z}) \to H_1(M_2, \mathbb{Z})$ induced by the inclusion map $j : M_1 \to M_2$ is an isomorphism, $p_Z = q$ and we are done.

Assume that the lemma holds when $\chi(K) \geq n$, $n \leq 0$, and suppose that $\chi(K) = n - 1$. Since $\chi(K) < 0$, there exist an analytic Jordan arc $\beta \in K \cap M_2^\circ$ with endpoints P_1, P_2 in $\partial(K) \cap \partial(M_1)$ and otherwise disjoint from $\partial(K)$, and such that $K - \beta$ contains no connected components with compact closure that are disjoint from $\partial(M_2)$. Let W be a trivial annular extension of $M_1 \cup \beta$ in M_2°, and notice that M_2 is an extension of W and $\chi(K_1) = \chi(K) + 1 \geq n$, where $K_1 := M_2 - W^\circ$. Consider an smooth loop $\gamma \subset W$ containing β and such that $H_1(W, \mathbb{Z}) = H_1(M_1 \cup \beta, \mathbb{Z}) = H_1(M_1, \mathbb{Z}) \oplus \mathbb{Z}[\gamma]$. Construct $X'_\sigma \in \mathcal{E}^*_N(M_1 \cup \beta)$ such that $X'|_{M_1} = X$ and $p_{X'_\sigma}(\gamma) = q(\gamma)$, and observe that $p_{X'_\sigma} = q|_{H_1(M_1 \cup \beta, \mathbb{Z})}$. Take a trivial annular extension W^* of W in M_2°, call R as a conformal compactification of W^* (hence of W), and write $E \subset R - W^*$ for a finite subset intersecting any component of $R - W^*$ at a unique point. By Theorem 4.1, there is $Y \in \mathcal{E}(R - E)$ such that $\|Y|_{M_1 \cup \beta} - X'|_{M_1 \cup \beta}\|_0 \leq \epsilon/2$ and $p_Y|_{H_1(M_1 \cup \beta, \mathbb{Z})} = p_{X'_\sigma}$.

Since $Y|_W \in \mathcal{E}_N(W)$, we can apply the induction hypothesis to W, M_2, and $Y|_W$ to find $Z \in \mathcal{E}_N(M_2)$ so that $\|Z|_W - Y|_W\|_0 \leq \epsilon/2$, $d_Z(W, \partial(M_2)) \geq C$ and $p_Z = q$. Therefore, $\|Z|_{M_1} - X\|_0 \leq \epsilon$ and $d_Z(M_1, \partial(M_2)) \geq d_Z(W, \partial(M_2)) \geq C$, closing the induction and proving the lemma. $\Box$

Theorem 4.2 (General Approximation) *Let M be a Riemann surface of finite conformal type and $\partial(M) \neq \emptyset$, and let M^* be an extension of M with $\partial(M^*) = \emptyset$. Consider an immersion $X \in \mathcal{E}_{M^*}(M)$ and a linear extension $q : H_1(M^*, \mathbb{Z}) \to \mathbb{R}$ of p_X.*

Then, there exists a sequence of conformal complete minimal immersions $\{Y_n\}_{n \in \mathbb{N}} \in \mathcal{E}(M^*)$ such that $\{Y_n|_M\}_{n \in \mathbb{N}} \to Y$ in the $\mathcal{C}^0(M)$-topology and $p_{Y_n} = q$.

Proof: It suffices to prove that for any $\epsilon > 0$ there is $Y_\epsilon \in \mathcal{E}(M^*)$ such that $\|Y_\epsilon|_M - X\|_0 \leq \epsilon$ and $p_Y = q$.

Let E^* denote the set of annular ends of finite conformal type in M^*, and denote by $\hat{M}^* := M^* \cup E^*$ the Riemann surface obtained by filling out the punctures in E^*. If $\hat{M}^*$ is compact, the theorem follows from Lemma 4.4, hence we can suppose that $\hat{M}^*$ is not compact. We will assume that $M^c \subset \hat{M}^*$ and write $E_0 = M^c \cap E^*$ (hence $M = M^c - E_0$).

Consider an exhaustion $\hat{N}_0 = M^c \subset \hat{N}_1 \subset \hat{N}_2 \subset \dots$ of $\hat{M}^*$ by compact regions such that $\hat{N}_{j+1}$ is an extension of $\hat{N}_j$ and $E_j := E^* \cap \hat{N}_j \subset \hat{N}_j^\circ$ for all $j \geq 0$. Write N_j for $\hat{N}_j - E_j$ and notice that $\hat{N}_j = N_j^c$, $j \geq 0$ (so $N_0 = M$). Set $Y_0 = X$ and use Lemma 4.4 to construct immersions $Y_j \in \mathcal{E}_{M^*}(N_j)$, $j \geq 1$, satisfying that:

(i) $\|Y_{j+1}|_{N_j} - Y_j\|_0 \leq \epsilon/2^{j+1}$ and $p_{Y_j} = q|_{H_1(N_j, \mathbb{R})}$, for all $j \geq 0$.

(ii) The intrinsic distance $d_{Y_{j+1}}\big(Y_{j+1}(N_j), Y_{j+1}\big(\partial(N_{j+1})\big)\big) \geq 1$ for all $j \geq 0$.

Let $Y_\epsilon : M^* \to \mathbb{R}^3$ be the possibly branched minimal immersion given by $Y|_{N_j} = \lim_{m \to \infty} Y_m|_{N_j}$, $j \in \mathbb{N}$, and note that $\lim_{m \to \infty} \|(Y_m - Y_\epsilon)|_{N_j}\|_0 = 0$ for all j and $\|Y_\epsilon|_M - X\|_0 \leq \epsilon$.

Let us show that Y_ϵ has no branch points. Indeed, let (g_m, ϕ_3^m) denote the Weierstrass data of Y_m, $m \in \mathbb{N}$, and likewise call (g, ϕ_3) as the ones of Y_ϵ. Obviously, $\{g_m, \phi_3^m)\}_{m \in \mathbb{N}} \to (g, \phi_3)$ uniformly on compact subsets of M^*. Take an arbitrary $p_0 \in M^*$, and consider $j_0 \in \mathbb{N}$ such that $p_0 \in N_{j_0}^\circ$. Up to a rigid motion, $g(p_0) \neq 0, \infty$, hence we can find an closed disc $D \subset N_{j_0}$ such that $p_0 \in D^\circ$ and $g_m|_D$, $m \in \mathbb{N}$, $g|_D$ are holomorphic and never vanishing. Since Y_m has no branch points, ϕ_3^m has no zeroes on D for all m. By Hurwith theorem, either $\phi_3 = 0$ of ϕ_3 has no zeroes on D as well. In the first case the identity principle gives $\phi_3 = 0$ on M^*, contradicting that $\|Y - X\|_0 < \epsilon$ provided that ϵ is small enough. This shows that Y_ϵ is a conformal minimal immersion.

Finally, let us see that Y_ϵ is complete and of WFTC. By Osserman's theorem, the Gauss map of Y_j extends meromorphically to $\hat{N}_j$, $j \in \mathbb{N}$. Since $\|Y_j - Y_\epsilon|_{N_j}\|_0$ is finite then $Y_j - Y_\epsilon$ extends harmonically to $\hat{N}_j$ for all $j \in \mathbb{N}$. So, the Gauss map of Y_ϵ extends meromorphically to $\hat{M}^*$ as well and $Y_\epsilon|_{N_j}$ has finite total curvature for any j. It remains to check that Y_ϵ is complete. Indeed, obviously those curves in M^* diverging to an end of E^* have infinite intrinsic length with respect to Y_ϵ. Moreover, condition (ii) implies that any curve in M^* diverging in $\hat{M}^*$ has also infinite intrinsic length and $Y_\epsilon \in \mathcal{E}(M^*)$. Clearly $p_{Y_\epsilon} = q$ and we are done. $\square$

Given an open Riemann surface N, an immersion $X \in \mathcal{E}(N)$ and $\theta \in \partial(\mathbb{D})$, we set $X_\theta = \operatorname{Re}\big(\int \theta \cdot \partial_z X\big)$ and call $\{X_\theta \ : \ \theta \in \partial(\mathbb{D})\} \subset \mathcal{E}(N)$ as the family of *associated minimal immersions* of X (well defined only when $\partial_z X$ is exact). The next corollary generalizes Pirola's results in [16]:

Corollary 4.2 *For any open Riemann surface N, there exists $Y \in \mathcal{E}(N)$ such that all its associated immersions are well defined. In particular, the space $\mathcal{E}(N) \neq \emptyset$.*

Proof: Fix a closed disc $D \subset N$ and an immersion $X \in \mathcal{E}_N(D)$. By Theorem 4.2, there is $\{Y_n\}_{n\in\mathbb{N}} \subset \mathcal{E}(N)$ such that $\{Y_n|_D\}_{n\in\mathbb{N}} \to X$ in the $\mathcal{C}^0(D)$-topology and $p_{Y_n} = q_0$ for all n, where $q_0 : H_1(N,\mathbb{Z}) \to \mathbb{R}$ is the null homomorphism. The corollary follows straightforwardly. $\square$

References

[1] A. Alarcón, F. J. López, *Minimal surfaces in $\mathbb{R}^3$ properly projecting into $\mathbb{R}^2$*. Preprint.

[2] A. Alarcon, L. Ferrer, F. Martin, *Density theorems for complete minimal surfaces in $\mathbb{R}^3$*. GAFA, **18 (1)** (2008), 1-49.

[3] E. Bishop, *Subalgebras of functions on a Riemann surface.* Pacific J. Math **8** (1958), 29-50.

[4] H. M. Farkas, I. Kra, *Riemann surfaces.* Graduate Texts in Math., **72**, Springer Verlag, Berlin, 1980.

[5] L. Ferrer, F. Martin, W. H. Meeks III, *The existence of proper minimal surfaces of arbitrary topological type.* Preprint.

[6] A. Huber, *On subharmonic functions and differential geometry in the large.* Comment. Math. Helv., **32**, 13-72 (1957).

[7] L. P. M. Jorge, W. H. Meeks III: *The topology of complete minimal surfaces of finite total Gaussian curvature.* Topology, Vol. **2** (1983), 203-221. 203-221.

[8] L.P. M. Jorge, F. Xavier, *A complete minimal surface in $\mathbb{R}^3$ between two parallel planes.* Ann. of Math., **112** (1980), 203-206.

[9] N. Kapouleas, *Complete embedded minimal surfaces of finite total curvature.* J. Differential Geom., **47** (1997), no. 1, 95-169.

[10] R. Kusner, N. Schmidt, *The spinor representation of minimal surfaces.* GANG Preprint III.27, 1993. Revised version, GANG Preprint 4.18 (arXiv:dg-ga/9512003v1).

[11] F. J. Lopez, *Exotic minimal surfaces.* In preparation.

[12] F. J. Lopez, *Uniform Approximation by nonorientable algebraic minimal surfaces in $\mathbb{R}^3$*. In preparation.

[13] F. J. Lopez, F. Martin and S. Morales, *Adding handles to Nadirashvili's surfaces.* J. Differ. Geom., **60**(1), 155-175 (2002).

[14] S. Morales, *On the existence of a proper minimal surface in R^3 with the conformal type of a disk.* GAFA, **13** (2003), 1281-1301.

[15] N. Nadirashvili, *Hadamard's and Calabi-Yau's conjectures on negatively curved and minimal surfaces.* Invent. math., **126** (1996), 457-465.

[16] G.P. Pirola, *Algebraic curves and non-rigid minimal surfaces in the Euclidean space.* Pacific J. Math. **183** (1998), 333-357.

[17] H. L. Royden, *Function Theory on Compact Riemann Surfaces.* J. Analyse Math., **18** (1967), 295-327.

[18] S. Scheinberg, *Uniform approximation by functions analytic on a Riemann surface.* Ann. of Math., **108** (1978), 257-298.

[19] S. Scheinberg, *Uniform approximation by meromorphic functions having prescribed poles.* Math. Ann., **243** (1979), 83-93.

[20] R. Osserman, *A survey of minimal surfaces.* Dover Publications, New York, second edition, 1986.

[21] B. White, *The bridge principle for stable minimal surfaces.* Calculus of Variations and P. D. E., **2** (1994), 405-425.

[22] B. White, *The bridge principle for unstable and for singular minimal surfaces.* Comm. Ann. Geom., **2**, No. 4 (1994), 513-532.

[23] S.-D. Yang, *A connected sum construction for complete minimal surfaces of finite total curvature.* Comm. Anal. Geom., **9** (2001), no. 1, 115-167.

FRANCISCO J. LOPEZ
Departamento de Geometría y Topología
Facultad de Ciencias, Universidad de Granada
18071 - GRANADA (SPAIN)
e-mail: fjlopez@ugr.es